


# Van Allen Probes reveal long-term behavior of Earth's ring current

May 19 2016, by Geoffrey Brown

---



During periods when there are no geomagnetic storms affecting the area around Earth (left image), high-energy protons (with energy of hundreds of thousands of electronvolts, or keV; shown here in orange) carry a substantial electrical current that encircles the planet, also known as the ring current. During periods when geomagnetic storms affect Earth (right), new low-energy protons (with energy of tens of thousands of electronvolts, or keV; shown here in magenta) enter the near-Earth region, enhancing the pre-existing ring current. Credit: Johns Hopkins APL

New findings based on a year's worth of observations from NASA's Van Allen Probes have revealed that the ring current - an electrical current carried by energetic ions that encircles our planet - behaves in a much different way than previously understood.

The ring current has long been thought to wax and wane over time, but the new observations show that this is true of only some of the particles, while other particles are present consistently. Using data gathered by the Radiation Belt Storm Probes Ion Composition Experiment, or RBSPICE, on one of the Van Allen Probes, researchers have determined that the high-energy protons in the ring current change in a completely different way from the current's low-energy protons. Such information can help adjust our understanding and models of the ring current - which is a key part of the space environment around Earth that can affect our satellites.

The findings were published in *Geophysical Research Letters*.

"We study the ring current because, for one thing, it drives a global system of electrical currents both in space and on Earth's surface, which during intense [geomagnetic storms](#) can cause severe damages to our technological systems," said lead author of the study Matina Gkioulidou, a space physicist at the Johns Hopkins University Applied Physics Laboratory in Laurel, Maryland. "It also modifies the [magnetic field](#) in near-Earth space, which in turn controls the motion of the radiation belt particles that surround our planet. That means that understanding the dynamics of the ring current really matters in helping us understand how radiation belts evolve as well."



Graphical rendering of the two Van Allen Probes Credit: JHU/APL, NASA

The ring current lies at a distance of approximately 6,200 to 37,000 miles (10,000 to 60,000 km) from Earth. The ring current was hypothesized in the early 20th century to explain observed global decreases in the Earth's surface magnetic field, which can be measured by ground magnetometers. Such changes of the ground magnetic field are described by what's called the Sym-H index.

"Previously, the state of the ring current had been inferred from the variations of the Sym-H index, but as it turns out, those variations represent the dynamics of only the low-energy protons," said Gkioulidou. "When we looked at the high-energy proton data from the RBSPICE instrument, however, we saw that they were behaving in a very different way, and the two populations told very different stories about the ring current."

The Van Allen Probes, launched in 2012, offer scientists the first chance in recent history to continuously monitor the ring current with instruments that can observe ions with an extremely wide range of energies. The RBSPICE instrument has captured detailed data of all types of these energetic ions for several years. "We needed to have an instrument that measures the broad energy range of the particles that carry the ring current, within the ring current itself, for a long period of time," Gkioulidou said. A period of one year from one of the probes was used for the team's research.

"After looking at one year of continuous ion data it became clear to us that there is a substantial, persistent ring current around the Earth even during non-storm times, which is carried by high-energy protons. During geomagnetic storms, the enhancement of the ring current is due to new, low-energy protons entering the near-Earth region. So trying to predict the storm-time ring current enhancement while ignoring the substantial pre-existing current is like trying to describe an elephant after seeing only its feet," Gkioulidou said.

Provided by NASA's Goddard Space Flight Center

Citation: Van Allen Probes reveal long-term behavior of Earth's ring current (2016, May 19)  
retrieved 10 April 2024 from

<https://phys.org/news/2016-05-van-allen-probes-reveal-long-term.html>

|                                                                                                                                                                                                                                          |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|