


Bose-Einstein condensates miscibility properties reveal surprises

June 27 2016



(a) Schematic illustration of the system. (b) Experimental result. The time evolution of the condensates shows clear bouncing, which are seemingly incompatible with the property of static miscible BECs, in which the ground state is spatially overlapped BECs. Credit: University of Electro Communications

Bose-Einstein condensates (BECs) are macroscopic systems that have quantum behaviour, and are useful for exploring fundamental physics. Now researchers at the Gakushuin University and the University of Electro-Communications have studied how the miscibility of multicomponent BECs affects their behaviour, with surprising results.

Fundamental particles have a property associated with angular momentum described as spin. Force particles - photons, gluons, and so on - have integer spin values and are called bosons; matter particles - electrons, neutrons, protons, and so on - have half integer values of spin and are called fermions. In composites of several fermions, as in atoms and nuclei, the total spin can be integer values so they can behave as bosons. While identical fermions cannot occupy the same state, bosons can, and if cooled to sufficiently low temperatures they will all occupy the lowest possible energy state - a Bose-Einstein condensate.

The researchers studied a BEC of rubidium atoms exploiting the element's rich spin states. They created optical traps containing around 3×10^5 atoms in two different [spin](#) states, and applied magnetic-field gradient pulses to separate condensates with different spins. The miscibility of different components of a BEC is determined by the strength of interactions between and within the [atoms](#), which the researchers could tune to produce miscible and immiscible multicomponent BECs.

After removing the magnetic field they left the system to evolve before releasing from the trap and imaged the resulting condensate distribution. "The various counterintuitive effects such as mutual penetration in immiscible BECs, bouncing between miscible BECs, and domain formation in miscible BECs were observed," report the researchers. Numerical simulations of the system revealed further insights, showing that "the properties of penetration and bouncing can be tuned by slightly changing the atomic interaction strengths."

More information: Yujiro Eto et al. Bouncing motion and penetration dynamics in multicomponent Bose-Einstein condensates, *Physical Review A* (2016). [DOI: 10.1103/PhysRevA.93.033615](https://doi.org/10.1103/PhysRevA.93.033615)

Provided by University of Electro Communications

Citation: Bose-Einstein condensates miscibility properties reveal surprises (2016, June 27)
retrieved 24 April 2024 from
<https://phys.org/news/2016-06-bose-einstein-condensates-miscibility-properties-reveal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.