


# Coexistence of superconductivity and charge density waves observed

June 21 2016

---



Scanning electron microscopy in combination with EELS electron spectroscopy permits to visualise atomic positions of the individual atoms in the heterostructure: Superconducting regions of YBaCuO are identified by yttrium (blue) and copper (pink), the ferromagnetic layers by manganese (green) and lanthanum (red). Credit: MPI Stuttgart

Physicists at BESSY II of Helmholtz-Zentrum Berlin studied an artificial structure composed of alternating layers of ferromagnetic and superconducting materials. Charge density waves induced by the interfaces were found to extend deeply into the superconducting regions, indicating new ways to manipulate superconductivity. The results are now being published in *Nature Materials*.

High-Tc superconductors were discovered 30 years ago: A class of ceramic metal oxide materials was found to pass electrical current without energy losses. In contrast to conventional superconductors that

have to be cooled almost to absolute zero, this property appears already at comparably high temperatures. In prototypical yttrium barium copper oxide (YBaCuO), the transition temperature is 92 Kelvin (minus 181 degrees centigrade). Hence, liquid nitrogen suffices as coolant to reach the superconducting phase. The discovery of [high-temperature superconductivity](#) has started a quest for applications, which are being implemented now.

Until now, however, the microscopic mechanism of high-Tc superconductivity is still matter of debate.

## **Superconducting and ferromagnetic thin layers**

A team of scientists lead by Prof. Bernhard Keimer, MPI for Solid State Research, and Dr. Eugen Weschke, HZB, have now investigated an artificial layer system composed of alternating nanolayers of YBaCuO and a ferromagnetic material. The thicknesses of the YBaCuO layers varied between 10 nm and 50 nm.

## **Tiny collective modulations of valence electrons observed**

As interfaces often determine the properties of such heterostructures, physicists were particularly interested in their role for the present system. During his PhD work using resonant x-ray diffraction at BESSY II, Alex Frano could detect tiny collective modulations of [valence electrons](#) around Cu atoms in the YBaCuO layer. Data analysis revealed that the resulting charge density wave does not remain located close to the interface but extends across the whole layer. " This finding is quite a surprise, as previous studies revealed a strong tendency of superconductivity to suppress the formation of [charge density waves](#)", explains Frano.

## Coexistence of superconductivity and charge density wave

"Engineering artificial interfaces in heterostructures of ferromagnetic and superconducting layers allowed to stabilize [charge density](#) waves even in the presence of [superconductivity](#): YBaCuO remains superconducting, while the charges arrange in a periodic structure", explains Weschke, "exploring the details of this coexistence on a microscopic scale is a challenging task for future experiments." A most exciting perspective of the present results is paving the way to controlling the superconducting state itself.

**More information:** A. Frano et al, Long-range charge-density-wave proximity effect at cuprate/manganate interfaces, *Nature Materials* (2016). [DOI: 10.1038/nmat4682](https://doi.org/10.1038/nmat4682)

Provided by Helmholtz Association of German Research Centres

Citation: Coexistence of superconductivity and charge density waves observed (2016, June 21) retrieved 28 April 2024 from <https://phys.org/news/2016-06-coexistence-superconductivity-density.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.