

Ice age bison fossils shed light on early human migrations in North America

June 6 2016



The steppe bison had much larger horns than modern bison. Radiocarbon dating and DNA analysis of bison fossils enabled researchers to track the migration of Pleistocene steppe bison into an ice-free corridor that opened along the Rocky Mountains about 13,000 years ago. Credit: Government of Yukon

Scientists using evidence from bison fossils have determined when an ice-

free corridor opened up along the Rocky Mountains during the late Pleistocene. The corridor has been considered a potential route for human and animal migrations between the far north (Alaska and Yukon) and the rest of North America, but when and how it was used has long been uncertain.

The researchers combined radiocarbon dating and DNA analysis to track the movements of [bison](#) into the corridor, showing that it was fully open by about 13,000 years ago. Their findings, published June 6 in *Proceedings of the National Academy of Sciences*, indicate that the corridor could not account for the initial dispersal of humans south of the ice sheets, but could have been used for later movements of people and animals, both northward and southward.

In the 1970s, geological studies suggested that the corridor might have been the pathway for the first movement of humans southward from Alaska to colonize the rest of the Americas. More recent evidence, however, indicated that the Cordilleran and Laurentide ice sheets coalesced at the height of the last ice age, around 21,000 years ago, closing the corridor much earlier than any evidence of humans south of the ice sheets. The initial southward movement of people into the Americas more than 15,000 years ago now seems likely to have been via a Pacific coastal route, but the Rocky Mountains corridor has remained of interest as a potential route for later migrations.

"The opening of the corridor provided new opportunities for migration and the exchange of ideas between people living north and south of the ice sheets," said first author Peter Heintzman, a postdoctoral researcher at UC Santa Cruz who led the DNA analysis.

Previous work by coauthor Beth Shapiro, professor of ecology and evolutionary biology at UC Santa Cruz, had shown that the bison populations north and south of the ice sheets were genetically distinct by

the time the corridor opened. By analyzing bison fossils from within the corridor region, the researchers were able track the movement of northern bison southward into the corridor and southern bison northward.

"The radiocarbon dates told us how old the fossils were, but the key thing was the genetic analysis, because that told us when bison from the northern and southern populations were able to meet within the corridor," Heintzman said.



Grant Zazula of the Yukon Paleontology Program with the skull of a Pleistocene steppe bison. Credit: Government of Yukon

The results showed that the southern part of the corridor opened first, allowing southern bison to start moving northward as early as 13,400 years ago, before the corridor fully opened. Later, there was some movement of northern bison southward, with the two populations overlapping in the corridor by 13,000 years ago.

"Bison fossils are the most widespread Quaternary mammal in western North America and of interest because they survived the extinctions at the end of the Pleistocene, unlike most other North American large mammals," said coauthor Duane Froese of the University of Alberta. "We were able to sample bison fossils, largely from museum collections, including critical ones from central Alberta that dated to the initial opening of the corridor."

According to Shapiro, archeological evidence suggests that human migration within the corridor was mostly from south to north. Sites associated with the Clovis hunting culture and its distinctive fluted point technology were widespread south of the corridor around 13,000 years ago and decline in abundance from south to north within the corridor region. A Clovis site in Alaska has been dated to no earlier than 12,400 years ago.

"When the corridor opened, people were already living south of there. And because those people were bison hunters, we can assume they would have followed the bison as they moved north into the corridor," Shapiro said.

The steppe bison of the Pleistocene (*Bison priscus*) were much bigger than modern bison (*Bison bison*), she said. Before the corridor closed, prior to the last glacial maximum, they moved freely up and down between the ice-free regions in the north and grasslands south of the ice sheets. After the ice sheets coalesced, the population that was cut off to the south contracted, leaving one genetically distinct southern lineage.

The DNA analysis used in this study focused on mitochondrial DNA, which is easier to recover from fossils than the DNA in chromosomes, because each cell has thousands of copies of the relatively short mitochondrial DNA sequence. While Shapiro's lab led the DNA analyses, Froese's lab led the radiocarbon dating work.

Many of the fossils they analyzed came from collections at the Royal Alberta Museum in Edmonton and other institutions. "Thousands of steppe bison fossils are recovered in northern Canada every year," said coauthor Grant Zazula of the Government of Yukon Palaeontology Program in Whitehorse. "Most of these fossils are uncovered by mining or gravel pit operators and later made available to scientists for study. These results speak to the importance of collecting and preserving fossils in order to better understand our history."

More information: Bison phylogeography constrains dispersal and viability of the Ice Free Corridor in western Canada, *PNAS*, www.pnas.org/cgi/doi/10.1073/pnas.1601077113

Provided by University of California - Santa Cruz

Citation: Ice age bison fossils shed light on early human migrations in North America (2016, June 6) retrieved 24 May 2024 from <https://phys.org/news/2016-06-ice-age-bison-fossils-early.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--