


Illuminating detection of deep cancers

June 20 2016



A luciferin analog generating near-infrared bioluminescence achieves highly sensitive deep-tissue imaging Credit: Tokyo Institute of Technology


To see right through something is to know its true intent—the same could be said to apply to biological tissues, especially tumors.

Bioluminescence imaging with a firefly enzyme, called luciferase, and its substrate D-luciferin, is widely used to monitor biological processes. However, the emission wavelength of bioluminescence produced by D-luciferin limits the sensitivity of this technique. At 562 nm, this light does not effectively penetrate [biological tissues](#).

To overcome this limitation, a team of Tokyo Tech and the University of Electro-Communications (UEC) researchers developed a luciferin analog (a compound that resembles another in structure) that can produce bioluminescence with near-infrared wavelength and is applicable in animal experiments. This allows markedly higher target-detection sensitivity, even at very low concentrations.

A novel soluble luciferin analog

The UEC researchers had previously synthesized a novel luciferin analog, AkaLumine, by altering the chemical structure of D-luciferin. While the emission wavelength of bioluminescence produced by AkaLumine yielded high penetration, its insolubility hindered its use. The team moved beyond this to screen for water-soluble derivatives of AkaLumine, and discovered that one of them, AkaLumine hydrochloride (AkaLumine-HCl), was in fact soluble. The Tokyo Tech researchers evaluated these substrates and had provided proper information for directing it to be practical use in [animal experiments](#), making AkaLumine-HCL applicable for bioluminescence imaging of deep tissues.




Optical imaging of the biological tissue (4-mm/8mm-thick sliced beef) were placed on the wells with each substrate (left). Penetration efficiency of bioluminescence through the tissue (right). Credit: Tokyo Institute of Technology

AkaLumine-HCl emitted near-infrared bioluminescence at 677 nm when reacted with firefly luciferase (Figure 1), and had greatly improved tissue-penetration efficiency. In 4-mm or 8-mm slice of beef, AkaLumine-HCl bioluminescence showed penetration 5-fold and 8.3-fold higher than bioluminescence produced by D-Luciferin (Figure 2). Notably, achieving such a high sensitivity using D-luciferin would require a 60-fold higher concentration.

To further evaluate the performance of AkaLumine-HCl in a lung cancer mouse model, the researchers compared the bioluminescence signals from mouse lung [cancer](#) treated with AkaLumine-HCl, D-luciferin, and its superior counterpart, cyclic alkylaminoluciferin (CycLuc1). Remarkably, AkaLumine-HCl significantly increased detection sensitivity of lung tumors as compared with D-luciferin and CycLuc1 (Figure 3).

Immediate applicability

Owing to its superior properties that enable higher [sensitivity](#) and accuracy, AkaLumine-HCl has potential to become the preferred choice for bioluminescence imaging. Nonetheless, for now, the benefits that its discovery brings can already be reaped in bioluminescence imaging studies in small animal models.



A luciferin analog generating near-infrared bioluminescence achieves highly sensitive deep-tissue imaging. Credit: Tokyo Institute of Technology

The study is published in *Nature Communications*.

More information: *Nature Communications*, [DOI:](#)

[10.1038/ncommms11856](https://phys.org/news/2016-06-illuminating-deep-cancers.html)

Provided by Tokyo Institute of Technology

Citation: Illuminating detection of deep cancers (2016, June 20) retrieved 27 April 2024 from <https://phys.org/news/2016-06-illuminating-deep-cancers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.