

Meadowsweet – good indication of restored river success

June 14 2016



Timber was floated in rivers in the 18th and 19th centuries. Credit: Swedish University of Agricultural Sciences

During the timber-floating era in rivers and streams, tributaries were cleared from all obstacles. Now, efforts have been made to restore many river reaches after these interferences. In order to measure the success of

such restorations, three researchers at Umeå University in Sweden have studied how to use meadowsweet to examine restoration outcomes. These results are published in the journal *Ecohydrology*.

The Swedish timber-floating era is long since history, and since, huge effort has been made into restoring affected river reaches. To do so in the best possible manner, it is important to follow up on the results of the restorations. Evaluations often focus on fish and aquatic insects whereas riparian and aquatic plants are rarely included. If there are no riparian plants, life for aquatic species is made very difficult. Therefore, riparian plants should be used as a measurement of the well-being of the watercourses.

"Using plants as a measurement of the health of nature, the so-called phytometer approach, is a rather unused method. What happens is that we breed a number of small seedlings and plant them in a variety of places in nature. After that, we follow their development. If the speed of their growth varies, it suggests that certain areas have different properties," says Anna Dietrich.

Anna Dietrich and her colleagues planted young seedlings of meadowsweet and purple moor grass on riparian zones of river reaches affected by timber-floating as well as restored rivers. They then compared how the plants survived and grew.

"Meadowsweet seedlings grew better on [riparian areas](#) by restored rivers and became both higher and sturdier than seedlings planted in channelized river reaches. Growth of stems and leaves was positively correlated to the restoration, which led to higher water levels in the summer. Longer floods in the spring also benefited the growth of roots, probably because the plants needed strong roots to hold on or else be flushed away," says Anna Dietrich.

Purple moor grass did not respond as clearly to restorations as the meadowsweet did apart from the fact that it grew better on restored rather than channelized riparian zones during the end of season.

Anna Dietrich took her doctorate at Umeå University in 2013 on the subject of phytometers, i.e. plants that can measure environmental changes. This is the publication of her fourth and last dissertation article.

About channelization and restoration:

When the Swedish forest industry expanded in the 19th and 20th centuries, tonnes of timber was shipped in rivers. Boulders, rocks and other obstacles made timber-floating cumbersome, which was solved by clearing and straightening the rivers to make them more similar to canals. The channels became homogenous, the water flow rapid and floods short but intense. Riparian zones suffered the consequences. In some parts, they were worn and torn by the rapid waterflow and by the actual timber. In other places they were left to dry behind man-made stone barriers. Plants, fish and other organisms were affected by the changes and some species even disappeared completely from these waters as a result of the river clearings that were made to facilitate timber-floating.

When restoring a watercourse, you create a more complex nature by returning boulders and gravel in watercourse and open up side channels again. That results in slower water flows, the water stays for longer in higher parts of watercourses where the number of floods in summer also increases. Water conditions are hence more similar to those of unspoiled watercourses. The quality of the soil is also improved after restoration. Based upon these facts, there is good hope that [plants](#) and animals will be able to re-establish. Restoration of watercourses after timber-floating is therefore highly ranked by conservationist authorities and organisations.

More information: Anna L. Dietrich et al. A phytometer study evaluating the effects of stream restoration on riparian vegetation, *Ecohydrology* (2016). [DOI: 10.1002/eco.1663](https://doi.org/10.1002/eco.1663)

Provided by Umea University

Citation: Meadowsweet – good indication of restored river success (2016, June 14) retrieved 19 April 2024 from <https://phys.org/news/2016-06-meadowsweet-good-indication-river-success.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.