

Microbial community dynamics dominate greenhouse gas production in thawing permafrost

June 22 2016



An international team examined microbial community dynamics at a site in Sweden where portions of the site are frozen permafrost and other areas are thawed (as shown here). They found that a particular microbe dominates the recently thawed areas, providing insights to improve climate predictions. Credit: Udo Schröter (Flickr) via a Creative Commons License

Northern permafrost ecosystems are changing rapidly, with rising temperatures causing many previously frozen environments to become wetlands such as fens. As permafrost thaws, the trapped organic carbon is accessible to microbial decomposition, which can release greenhouse gases (GHGs) to the atmosphere. A team led by the University of Arizona examined microbial community dynamics at a site in Sweden where portions of the site are frozen permafrost and other areas are thawed. They found that a single microbe dominated recently thawed sites, with its relative abundance strongly correlating with the magnitude and specific type of methane produced at any given site.

The results represent a significant advance in understanding and accurately modeling the critical biogeochemical processes of microbes in permafrost. The findings will improve predictions of climate change impacts on these delicate ecosystems and of potential consequences to the atmosphere. Finding the striking dominance of a single microbial species in mediating the large-scale carbon cycle process is highly unusual and provides an opportunity to more effectively track and predict the impacts of [climate change](#) across an entire region. Incorporating the datasets will facilitate development of ecosystem-scale models of the processes.

Understanding of [microbial communities](#) that can release [greenhouse gases](#) (GHGs) is limited, especially the specific nature of processes that affect rates of carbon decomposition and the balance of released carbon

dioxide (CO₂) versus methane (CH₄). Because CH₄ is a much more potent GHG in the short term, understanding the microbial mechanisms driving the large-scale processes of both gases would significantly improve the ability to predict possible [climate change impacts](#). An international, interdisciplinary team of researchers led by the University of Arizona has examined microbial community dynamics at a site in northern Sweden located in a natural temperature gradient. Northern portions of this site are frozen permafrost, while southern areas are thawed wetland fens.

Over several years, the team (1) measured CO₂ and CH₄ production along the gradient, (2) examined isotopic signatures of the gases characteristic of distinct microbial processes, and (3) correlated the data with measured shifts in microbial community composition and abundance. Only small amounts of GHGs were released from frozen permafrost, but progressively more were released, predominantly CH₄, with increasing temperatures and thawing. The team linked these observations with extensive shifts in microbial community composition, revealing a reproducible succession pattern of different types of CH₄-producing microbes (methanogens) across the thaw gradient. Surprisingly, a single methanogen species, *Candidatus Methanoflorens stordalenmirensis*, was dominant in recently thawed sites, with its relative abundance strongly correlating with the magnitude and specific type of CH₄ produced at any given site.

More information: Carmody K. McCalley et al. Methane dynamics regulated by microbial community response to permafrost thaw, *Nature* (2014). [DOI: 10.1038/nature13798](https://doi.org/10.1038/nature13798)

Provided by US Department of Energy

Citation: Microbial community dynamics dominate greenhouse gas production in thawing permafrost (2016, June 22) retrieved 23 April 2024 from <https://phys.org/news/2016-06-microbial-dynamics-dominate-greenhouse-gas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.