


New radio map of Jupiter reveals what's beneath colorful clouds

June 2 2016



In this animated gif, optical images of the surface clouds encircling Jupiter's equator --including the famous Great Red Spot -- alternate with new detailed radio images of the deep atmosphere (up to 30 kilometers below the clouds). The radio map shows ammonia-rich gases rising to the surface (dark) intermixed with descending, ammonia-poor gases (bright). In the cold temperatures of the upper atmosphere (160 to 200 Kelvin, or -170 to -100 degrees Fahrenheit), the rising ammonia condenses into clouds, which are invisible in the radio region. Credit: Radio: Robert J. Sault (Univ. Melbourne), Imke de Pater and Michael H. Wong (UC Berkeley). Optical: Marco Vedovato, Christopher Go, Manos Kardasis, Ian Sharp, Imke de Pater.

Astronomers using the upgraded Karl G. Jansky Very Large Array in

New Mexico have produced the most detailed radio map yet of the atmosphere of Jupiter, revealing the massive movement of ammonia gas that underlies the colorful bands, spots and whirling clouds visible to the naked eye.

The University of California, Berkeley researchers measured radio emissions from Jupiter's atmosphere in wavelength bands where clouds are transparent. The observers were able to see as deep as 100 kilometers (60 miles) below the cloud tops, a largely unexplored region where clouds form.

The planet's thermal [radio emissions](#) are partially absorbed by ammonia gas. Based on the amount of absorption, the researchers could determine how much ammonia is present and at what depth.


By studying these regions of the planet's atmosphere, astronomers hope to learn how global circulation and cloud formation are driven by Jupiter's powerful internal heat source. These studies also will shed light on similar processes occurring on other giant planets in our solar system and on newly discovered giant exoplanets around distant stars.

"We in essence created a three-dimensional picture of ammonia gas in Jupiter's atmosphere, which reveals upward and downward motions within the turbulent atmosphere," said principal author Imke de Pater, a UC Berkeley professor of astronomy.

The map bears a striking resemblance to visible-light images taken by amateur astronomers and the Hubble Space Telescope, she said.

The radio map shows ammonia-rich gases rising into and forming the upper cloud layers: an ammonium hydrosulfide cloud at a temperature near 200 Kelvin (minus 100 degrees Fahrenheit) and an ammonia-ice cloud in the approximately 160 Kelvin cold air (minus 170 degrees

Fahrenheit). These clouds are easily seen from Earth by optical telescopes.



The VLA radio map of the region around the Great Red Spot in Jupiter's atmosphere shows complex upwellings and downwellings of ammonia gas (upper map), that shape the colorful cloud layers seen in the approximately true-color Hubble map (lower map). Two radio wavelengths are shown in blue (2 cm) and gold (3 cm), probing depths of 30-90 kilometers below the clouds. Credit: Radio:

Michael H. Wong, Imke de Pater (UC Berkeley), Robert J. Sault (Univ. Melbourne). Optical: NASA, ESA, A.A. Simon (GSFC), M.H. Wong (UC Berkeley), and G.S. Orton (JPL-Caltech)

Conversely, the radio maps show ammonia-poor air sinking into the planet, similar to how dry air descends from above the [cloud layers](#) on Earth.

The map also shows that hotspots—so-called because they appear bright in radio and thermal infrared images—are ammonia-poor regions that encircle the planet like a belt just north of the equator. Between these hotspots are ammonia-rich upwellings that bring ammonia from deeper in the planet.

"With radio, we can peer through the clouds and see that those hotspots are interleaved with plumes of ammonia rising from deep in the planet, tracing the vertical undulations of an equatorial wave system," said UC Berkeley research astronomer Michael Wong.

The final maps have the best spatial resolution ever achieved in a radio map: 1,300 kilometers.

"We now see high ammonia levels like those detected by Galileo from over 100 kilometers deep, where the pressure is about eight times Earth's atmospheric pressure, all the way up to the cloud condensation levels," de Pater said.

De Pater, Wong and their colleagues will report their findings and highly detailed maps in the June 3, 2016 issue of the journal *Science*.

Prelude to Juno's arrival

The observations are being reported just one month before the July 4, 2016 arrival at Jupiter of NASA's Juno spacecraft, which plans, in part, to measure the amount of water in the deep atmosphere where the Very Large Array looked for ammonia.

"Maps like ours can help put their data into the bigger picture of what's happening in Jupiter's atmosphere," de Pater said, noting that her team will observe Jupiter with the VLA at the same time as Juno's microwave instruments are probing for water.

Key to the new observations was an upgrade to the VLA that improved sensitivity by a factor of 10, said Bryan Butler, a co-author and staff astronomer at the National Radio Astronomy Observatory in Socorro, New Mexico, which operates the VLA. "These Jupiter maps really show the power of the upgrades to the VLA."

The team observed over the entire frequency range between 4 and 18 gigahertz (1.7—7 centimeter wavelength), which enabled them to carefully model the atmosphere, said David DeBoer, a research astronomer with UC Berkeley's Radio Astronomy Laboratory.

"We now see fine structure in the 12 to 18 gigahertz band, much like we see in the visible, especially near the Great Red Spot, where we see a lot of little curly features," Wong said. "Those trace really complex upwelling and downwelling motions there."

The observations also resolve a puzzling discrepancy between the ammonia concentration detected by the Galileo probe when it plunged through the atmosphere in 1995—4.5 times the abundance observed in the sun—and VLA measurements from before 2004, which showed much less [ammonia gas](#) than measured by the probe.

"Jupiter's rotation once every 10 hours usually blurs radio maps, because

these maps take many hours to observe," said co-author Robert Sault, of the University of Melbourne in Australia. "But we have developed a technique to prevent this and so avoid confusing together the upwelling and downwelling ammonia flows, which had led to the earlier underestimate."

More information: "Peering through Jupiter's clouds with radio spectral imaging," *Science*, [DOI: 10.1126/science.aaf2210](https://doi.org/10.1126/science.aaf2210)

Provided by University of California - Berkeley

Citation: New radio map of Jupiter reveals what's beneath colorful clouds (2016, June 2) retrieved 24 April 2024 from <https://phys.org/news/2016-06-radio-jupiter-reveals-beneath-clouds.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.