


Universe's first life might have been born on carbon planets

June 7 2016



In this artist's conception, a carbon planet orbits a sunlike star in the early universe. Young planetary systems lacking heavy chemical elements but relatively rich in carbon could form worlds made of graphite, carbides and diamond rather than Earth-like silicate rocks. Blue patches show where water has pooled on the planet's surface, forming potential habitats for alien life. Christine

Pulliam (CfA). Sun image: NASA/SDO

Our Earth consists of silicate rocks and an iron core with a thin veneer of water and life. But the first potentially habitable worlds to form might have been very different. New research suggests that planet formation in the early universe might have created carbon planets consisting of graphite, carbides, and diamond. Astronomers might find these diamond worlds by searching a rare class of stars.

"This work shows that even [stars](#) with a tiny fraction of the [carbon](#) in our solar system can host planets," says lead author and Harvard University graduate student Natalie Mashian.

"We have good reason to believe that alien life will be carbon-based, like life on Earth, so this also bodes well for the possibility of life in the [early universe](#)," she adds.

The primordial universe consisted mostly of hydrogen and helium, and lacked chemical elements like carbon and oxygen necessary for life as we know it. Only after the first stars exploded as supernovae and seeded the second generation did planet formation and life become possible.

Mashian and her PhD thesis advisor Avi Loeb (Harvard-Smithsonian Center for Astrophysics) examined a particular class of old stars known as carbon-enhanced metal-poor stars, or CEMP stars. These anemic stars contain only one hundred-thousandth as much iron as our Sun, meaning they formed before interstellar space had been widely seeded with heavy elements.

"These stars are fossils from the young universe," explains Loeb. "By studying them, we can look at how planets, and possibly [life](#) in the

universe, got started."

Although lacking in iron and other [heavy elements](#) compared to our Sun, CEMP stars have more carbon than would be expected given their age. This relative abundance would influence [planet formation](#) as fluffy carbon dust grains clump together to form tar-black worlds.

From a distance, these carbon planets would be difficult to tell apart from more Earth-like worlds. Their masses and physical sizes would be similar. Astronomers would have to examine their atmospheres for signs of their true nature. Gases like carbon monoxide and methane would envelop these unusual worlds.

Mashian and Loeb argue that a dedicated search for planets around CEMP stars can be done using the transit technique. "This is a practical method for finding out how early planets may have formed in the infant universe," says Loeb.

"We'll never know if they exist unless we look," adds Mashian.

This research has been accepted for publication in the *Monthly Notices of the Royal Astronomical Society* and is available [online](#).

More information: CEMP stars: possible hosts to carbon planets in the early universe, arxiv.org/abs/1603.06943

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Universe's first life might have been born on carbon planets (2016, June 7) retrieved 19 April 2024 from <https://phys.org/news/2016-06-universe-life-born-carbon-planets.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.