

# Analysis of anatomy and diet finds evolution follows least resistant path

July 1 2016

---

Evolution follows the path of least resistance, which can result in suboptimal physical traits that don't ideally match the functional need, according to a new analysis by University of Arkansas anthropologist Peter Ungar.

Ungar, distinguished professor, chair of Department of Anthropology and director of the environmental dynamics doctoral program at the University of Arkansas, detailed his findings in *Science* this month with coauthor Leslea Hlusko, associate professor at the University of California Berkeley.

"Paleontologists typically reconstruct past behavior by assuming that function follows form," Ungar said. "We need to look at things in a different way and consider the number of genetic steps it takes to get from one anatomy to another. There can be more than one function for a given form and different forms can serve the same function."

Ungar analyzed teeth from two human ancestors with similar dental and jaw structure and found that dental anatomy historically associated with a hard food diet was used to eat mainly plant-based diets. And while the two species had similar anatomies and teeth, their diets were different.

The *Paranthropus boisei* from eastern Africa and *Paranthropus robustus* from southern Africa are groups of species from 4.2 to 1.3 million years ago that share similar head and tooth anatomy characterized by large, flat teeth with thick enamel and jaws and faces that indicate strong chewing

muscles.

Those characteristics have often been interpreted as adaptive for crushing hard foods. Ungar and his colleagues analyzed the microwear on the teeth and used previously published data on carbon isotopes to conclude that *P. robustus* had a diet of plant foods and only occasional consumption of hard objects like nuts or roots, while the microwear and isotope values on the teeth of *P. boisei* indicate it consumed softer, tougher and possibly more abrasive foods like grasses.

"So despite their similar masticatory morphology, chemical and wear traces of the foods eaten suggest that these two species differed markedly in their diets," Ungar said.

And neither diet matched previous inferences that the large muscles and facial structure indicated a diet of hard, crunchy food.

More crested molars like those in gorillas would be more efficient in tearing fibrous plant material, but flatter molars with thicker enamel, like those of modern humans, advanced through the genetic record.

Ungar said the form of the tooth and jaw anatomy in this case is ideal for the diets these hominins ate, but it took fewer genetic changes to get there. Thicker enamel with a simpler tooth architecture was more adaptable for a variety of challenging diets through time, so it carried on as the human species developed.

"You will reach suboptimal, but functional, solutions if they require fewer genetic mutations," Ungar said. "We need to look at things in a new way and take into account the evolutionary path of least resistance."

**More information:** P. S. Ungar et al. The evolutionary path of least resistance, *Science* (2016). [DOI: 10.1126/science.aaf8398](https://doi.org/10.1126/science.aaf8398)

Provided by University of Arkansas

Citation: Analysis of anatomy and diet finds evolution follows least resistant path (2016, July 1)  
retrieved 19 April 2024 from

<https://phys.org/news/2016-07-analysis-anatomy-diet-evolution-resistant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.