

New approach for environmental test on livestock drugs

July 27 2016



Dung pat with a specific concentration of ivermectin. Credit: UZH

Drugs for livestock can harm beneficial organisms that break down

dung. Therefore newly developed medical substances need to be tested on single species in the lab. An international research group including evolutionary biologists from the University of Zurich have been scrutinizing the reliability of such laboratory tests, evaluating the implementation of a field test based on the anti-parasitic drug ivermectin at four climatically different locations. The scientists thus presented a novel approach for more advanced environmental compatibility tests.

Livestock medications can impair beneficial organisms that break down dung. Too high a dosage of ivermectin, a common drug against parasites, harms coprophilous organisms, for instance. The toxicity of new livestock medications therefore needs to be verified in ecotoxicological tests with individual animal species such as the common yellow dung fly, the barn fly or a dung beetle. This involves determining the lethal dose leading to the death of half the maggots (LD50 test). However, sensitivity to toxic substances is known to vary significantly even among closely related coprophilous organisms, which begs the question as to how representative the reaction of any individual animal species actually is in such laboratory tests. After all, there is a high risk that more sensitive species will continue to be harmed by the substance, jeopardizing key ecosystem functions in the long run.

An international research group including UZH evolutionary biologist Wolf Blanckenhorn recently proposed extending the testing scheme to a representative selection of all organisms that break down dung, ideally in their natural environment. The scientists now presented a successful and more comprehensive higher-tier ecotoxicological field test. Their study provides important insights into minimizing the risks of drug residues in nature.

Earthworms compensate for loss of coprophilous insects

For their feasibility study, the scientists worked on cattle pastures in the Canadian Prairie and the agricultural landscapes of southern France, the Netherlands and Switzerland - four locations with very different climatic conditions. On these pastures, they distributed dung pats with different concentrations of ivermectin. "As expected, the overall number and diversity of [dung beetles](#), [dung flies](#) and parasitoid wasps decreased as the ivermectin concentration increased," explains Blanckenhorn. However, a number of species also proved to be resistant: earthworms and springtails living in the ground underneath the cowpats were not notably affected, and a parallel test ultimately revealed that dung degradation was not significantly impaired. "Evidently, [beneficial organisms](#) not affected as much by the drug, such as earthworms, were apparently able to compensate for the loss of other organisms," sums up Blanckenhorn.

A basis for decision makers and licensing authorities

Despite diverse environmental conditions and methodological details, the results were very similar and reproducible in all four habitats. "Our field approach was therefore a success and in principle can be recommended. The regulation authorities responsible, such as the European Medicines Agency EMA, now have to decide whether this more conclusive yet more complex test should be required in the future," says Blanckenhorn. The amount of effort involved in determining the numerous dung organisms is tremendous and impossible without expert biological knowledge. "Classifying species via so-called DNA barcoding, based on each organism's unique genetic fingerprint, is possible in principle and will probably be more cost-effective in the future. However, this approach requires the establishment of a complete database for coprophilous [organisms](#), which does not yet exist," concludes the scientist.

More information: Kevin D. Floate, Wolf U. Blanckenhorn. Special

Section: Non-target Structural and Functional Effects of Ivermectin Residues in Cattle Dung on Pasture – Guidance for Researchers and Regulators. *Environmental Toxicology and Chemistry*. Volume 35, Issue 8. July 21, 2016. [DOI: 10.1002/etc.3549](https://doi.org/10.1002/etc.3549)

Provided by University of Zurich

Citation: New approach for environmental test on livestock drugs (2016, July 27) retrieved 27 July 2024 from <https://phys.org/news/2016-07-approach-environmental-livestock-drugs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.