


Asymmetrical magnetic microbeads turned into micro-robots

July 20 2016



Credit: SciencePOD

Janus was a Roman god with two distinct faces. Thousands of years later, he inspired material scientists working on asymmetrical microscopic spheres—with both a magnetic and a non-magnetic half—called Janus particles. Instead of behaving like normal magnetic beads, with opposite poles attracting, Janus particle assemblies look as if poles of the same type attract each other.



A new study reveals that the dynamics of such assemblies can be predicted by modelling the interaction of only two particles and simply taking into account their magnetic asymmetry. These findings were recently published in *EPJ E* by Gabi Steinbach from the Chemnitz University of Technology, Germany, and colleagues at the Helmholtz-Zentrum Dresden-Rossendorf. It is part of a topical issue entitled "Nonequilibrium Collective Dynamics in Condensed and Biological Matter." The observed effects were exploited in a lab-on-a-chip application in which microscopic systems perform tasks in response to a changing external magnetic field.

Initially, the authors experimentally investigated large assemblies of these microbeads with an optical microscope. They then performed numerical simulations based on a simple model featuring off-centred magnetic dipoles, and performed a detailed comparison of both experimental and simulation approaches. Under an oscillating field, the particles continuously revert back to standard magnetic behaviour where the North magnetic poles of particles point towards other particles' South poles; and this behaviour has been confirmed for as few as two particles.

The authors concluded that the interactions between the off-centred dipoles with the changing external magnetic field transform larger particle clusters into different configurations. For example, they found that staggered bead chains continuously expand to spontaneously form linear chains. This could be harnessed, for instance, to create a zipper-style micro-muscle on a chip; here a staggered chain repeatedly expands and contracts depending on the intensity of the applied field.

More information: Gabi Steinbach et al. Non-equilibrium dynamics of magnetically anisotropic particles under oscillating fields, *The European Physical Journal E* (2016). DOI: 10.1140/epje/i2016-16069-6



Provided by Inderscience Publishers

Citation: Asymmetrical magnetic microbeads turned into micro-robots (2016, July 20) retrieved 20 April 2024 from

https://phys.org/news/2016-07-asymmetrical-magnetic-microbeads-micro-robots.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.