

# Microswimmer robot chains can decouple and reconnect in magnetic field

July 28 2016

---



Microswimmer robots, resembling a chain of beads, can be made to move through a liquid by rotating in a magnetic field. New research shows how they can split into separate operable robots and then reconnect. Credit: Drexel University

Drexel University researchers, led by MinJun Kim, PhD, a professor in the College of Engineering, have successfully pulled off a feat that both sci-fi fans and Michael Phelps could appreciate. Using a rotating

magnetic field they show how multiple chains of microscopic magnetic bead-based robots can link up to reach impressive speeds swimming through in a microfluidic environment. Their finding is the latest step toward using the so-called "microswimmers" to deliver medicine and perform surgery inside the body.

In a paper recently published in *Nature Scientific Reports*, the mechanical engineers describe their process for magnetically linking and unlinking the beads while they're swimming, and individually controlling the smaller decoupled robots in a [magnetic field](#). This data helps further the concept of using microrobots for targeted, intravenous [drug delivery](#), surgery and cancer treatment.

"We believe microswimmer robots could one day be used to carry out medical procedures and deliver more direct treatments to affected areas inside the body," said U Kei Cheang, PhD, a postdoctoral research fellow in Drexel's College of Engineering and lead author of the paper. "They can be highly effective for these jobs because they're able to navigate in many different biological environments, such as the blood stream and the microenvironment inside a tumor."

One of the central findings is that longer chains can swim faster than shorter ones. This was determined by starting with a three-bead swimmer and progressively assembling longer ones. The longest chain examined by the group, 13-beads in length, reached a speed of 17.85 microns/second.

Drexel engineers have been adding the understanding of microrobots for biomedical applications for nearly a decade, with the goal of producing a robotic chain that can travel inside the body, then decouple to deliver their medicinal payload or targeted treatment.

The reason for this approach is that a rather versatile robot that can do

multiple tasks could be controlled using a single magnetic field.

The robot chains move by spinning, like a long screw-like propeller in step with a rotating [external magnetic field](#). So the faster the field rotates, the more the robots spin and the faster they move. This dynamic propulsion system is also the key to getting them to divide into shorter segments. At a certain rate of rotation the robotic chain will split into two smaller chains that can move independently of each other.



By splitting the microswimmer robot chain into separate actuated robots, multiple tasks can be completed at once. This makes them ideal for surgery and targeted drug delivery. Credit: Drexel University

"To disassemble the microswimmer we simply increased the rotation frequency," Cheang said. "For a seven-bead microswimmer, we showed

that by upping the frequency 10-15 cycles the hydrodynamic stress on the swimmer physically deformed it by creating a twisting effect which lead to disassembly into a three-bead and four-bead swimmer."

Once separate, the field can be adjusted to manipulate the three and four-bead robots to move in different directions. Because the beads are magnetized, they can eventually be reconnected—simply by tweaking the field to bring them back into contact on the side with the corresponding magnetic charge. The team also determined optimal rotation rates and angle of approach to facilitate re-linking the microswimmer chains.



By rotating the magnetic field at a certain frequency the robotic chains will split

into separate, individually controllable robots. Credit: Drexel University

This finding is a key component of a larger project in which Drexel is partnering with 10 institutions of research and medicine from around the world to develop this technology for performing [minimally invasive surgery](#) on blocked arteries.

"For applications of drug delivery and minimally invasive surgery, future work remains to demonstrate the different assembled configurations can achieve navigation through various in vivo environments, and can be constructed to accomplish different tasks during operative procedures," the authors write. "But we believe that the mechanistic insight into the assembly process we discussed in this research will greatly aid future efforts at developing configurations capable of achieving these crucial abilities."

**More information:** *Scientific Reports*, [DOI: 10.1038/srep30472](https://doi.org/10.1038/srep30472) , [www.nature.com/articles/srep30472](http://www.nature.com/articles/srep30472)

Provided by Drexel University

Citation: Microswimmer robot chains can decouple and reconnect in magnetic field (2016, July 28) retrieved 3 May 2024 from <https://phys.org/news/2016-07-microswimmer-robot-chains-decouple-reconnect.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--