


NASA finds Tropical Cyclone Frank fading

July 28 2016



On July 27, the RapidScat instrument observed Frank's strongest sustained winds were near 30 meters per second (red) north of the center. Credit: NASA JPL, Alex Winteer

Tropical Storm Frank weakened to a tropical storm on July 28, 2016 and


satellite imagery shows that winds and thunderstorms have continued to weaken.

On July 27, NASA's RapidScat instrument observed Frank's strongest sustained winds were near 30 meters per second (67 mph/108 kph) north of the center. Tropical-storm-force winds extended outward up to 60 miles (95 km) from the center. Frank weakened from a hurricane to a [tropical storm](#) by 5 p.m. EDT on July 27.

The RapidScat instrument flies aboard the International Space Station and measures winds over an ocean surface. Images are created at NASA's Jet Propulsion Laboratory in Pasadena, California.

Later on July 27, at 21:35 UTC (5:35 p.m. EDT) the Atmospheric Infrared Sounder or AIRS instrument aboard NASA's Aqua satellite captured infrared data of Frank. The AIRS data showed that the coldest cloud top temperatures of at least minus 63 degrees Fahrenheit (minus 53 degrees Celsius) and strongest thunderstorms were only along the southern quadrant of the storm.

At 5 a.m. EDT (0900 UTC), the center of Tropical Storm Frank was located near 23.6 degrees latitude and 123.7 degrees west longitude. That's about 875 miles (1,410 km) west of the southern tip of Baja California. Maximum sustained winds have decreased to near 40 mph (65 kph) with higher gusts.



On July 27 at 21:35 UTC, the AIRS instrument aboard NASA's Aqua satellite captured an image of the strongest storms (purple) remaining in Frank, south of the center. Credit: NASA JPL, Ed Olsen


Frank was moving toward the west-northwest near 9 mph (15 kph), and the National Hurricane Center expects a gradual turn toward the west and west-southwest over the next 48 hours as the storm weakens.

NOAA GOES-West infrared image of Tropical Storm Frank was taken on July 28, 2016 at 1315 UTC (9:15 a.m. EDT). The GOES-West image showed very weak storms and mostly along the southern quadrant, as the

infrared imagery from AIRS had shown 16 hours before. The GOES image was generated at the NASA/NOAA GOES Project at NASA's Goddard Space Flight Center, Greenbelt, Maryland.

Forecaster Pasch at NOAA's National Hurricane Center (NHC) said that "All of the deep convection associated with Frank has dissipated, and since the cyclone is moving over waters cooler than 23 [degrees](#) Celsius, regeneration of thunderstorms within the circulation is not anticipated. Therefore, the cyclone is likely to degenerate into a post-tropical remnant low pressure area later today.

NHC said that the storm will continue to weaken as the circulation spins down over the next couple of days. Frank is expected to dissipate by Sunday, July 31.



This NOAA GOES-West infrared image of Tropical Storm Frank was taken on July 28, 2016 at 1315 UTC (9:15 a.m. EDT). Credit: NASA NOAA GOES Project

Provided by NASA's Goddard Space Flight Center

Citation: NASA finds Tropical Cyclone Frank fading (2016, July 28) retrieved 26 June 2024 from <https://phys.org/news/2016-07-nasa-tropical-cyclone-frank.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.