

Cavitands enhance selectivity and hydrolysis rate of one ester on a long-chain diester

August 11 2016, by Heather Zeiger

A challenge for chemists is trying to selectively react one of two identical functional groups on the same molecule. For example, a diester with a long carbon chain in between the two ester groups has two chemically equivalent sites. Hydrolysis of this diester results in a mixture of products where one or both of the esters are converted to carboxylic acids. This results in lower overall yields.

New research by Qixun Shi, Matthew P. Mower, Donna G. Blackmond, and Julius Rebek, Jr. of Fudan University and Scripps Research Institute describe the use of cavitands as a way to overcome the monofunctionalization problem. They found that cavitands force a long-chain α,ω -dimethyl ester to fold onto itself changing the molecule's behavior compared to bulk solution. The cavitand results in greater selectivity and [reaction](#) rates for both the acid and base hydrolysis of diesters. Their work appears in the *Proceedings of the National Academy of Sciences*.

While there are ways to functionalize one of two identical functional groups on a molecule, these methods rely on there being some difference between the two functional groups. But, in the case of multi-carbon diesters, these procedures do not work because the functional groups are so far apart from each other that they essentially behave as two reactants (i.e., they have equivalent rate constants). One possible solution is to entrap the long-chain reactant within a molecular container causing the reactant to fold in a particular fashion. This is similar to how the body entraps a peptide chain in a chaperone protein during protein synthesis

or how enzymes will catalyze an otherwise difficult reaction by causing a guest molecule to fold into a conformation that buries some of the reactive sites.

Cavitands are macrocyclic molecules shaped like a cup. The lip of the cup has functional groups that serve to broaden the rim and prevent dimerization. The chemical composition within the cavity as well as the cavity's shape promotes guest folding into a particular conformation. In the current research, the authors found that the α,ω -dimethyl ester guest folded into a J-shape, with one end outside of the cup while the other end remains within the cup. This allowed for both acid promoted and base-promoted (i.e., saponification) hydrolysis of only one end of the ester.

Using NMR to follow the reaction process, Shi, et al. first tested acid hydrolysis using DCl added to D₂O in a solution containing diester and the cavitand. One end of the ester is deep within the cavitand while the other is accessible for hydrolysis. Once one end is hydrolyzed the other ester is rendered less reactive; however, because the J-shape conformation is dynamic, some of the other end of the diester does hydrolyze. The yield for the monoester product is still significantly greater than in bulk solution based on rate data.

Kinetic studies using NMR peak integration of the acid hydrolysis reaction indicate that the ester portion of the molecule remains suppressed within the cavitand three-fourths of the time. The reaction rate is more than a factor of ten greater than the comparable reaction in bulk solution. The authors suggest that this may be due to better solvation of the exposed ester. With the organic molecule-cavitand complex, the reaction can be carried out in water. Without the cavitand, the acid hydrolysis reaction requires the addition of an organic co-solvent.

The saponification reaction demonstrated even better selectivity and enhanced reaction rates compared to the acid hydrolysis reaction. The major product was monester with

Citation: Cavitands enhance selectivity and hydrolysis rate of one ester on a long-chain diester (2016, August 11) retrieved 16 July 2024 from <https://phys.org/news/2016-08-cavitands-hydrolysis-ester-long-chain-diester.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.