


Using fast particles to probe hot matter in nuclear collisions

August 1 2016



Propagation of fast subatomic particles (quarks and gluons) through the hot and dense matter called quark-gluon plasma created in high-energy nuclear collisions. Lines with arrows indicate momenta of fast particles; colors indicate the local density of the quark-gluon plasma. The insert shows jet transport coefficient extracted by the JET Collaboration. Credit: Lawrence Berkeley National Laboratory

The hottest matter that existed in the early universe after the Big Bang is created in collisions of high-energy nuclei. Using information on the

propagation and attenuation of fast particles coming from the collisions, nuclear physicists can extract transport properties of the hot, dense matter.

The extracted jet transport coefficients, which have the highest precision to date, show how well the [quark-gluon plasma](#) (QGP) can transfer momentum and energy within its bulk. This result will lead to a much better understanding of the plasma constituents and properties.

Heavy nuclei are accelerated to nearly the speed of light and collide with each other to create a new state of matter called QGP that only existed in the [early universe](#) shortly after the Big Bang. The JET Topical Collaboration, funded by Department of Energy and lead by the Department of Energy's Lawrence Berkeley National Laboratory, investigated properties of QGP using information on fast particles, which are produced early in the collision and propagate through the plasma—much like a CAT scan in medical imaging. Combining theoretical modeling with data from experiments at the Relativistic Heavy-Ion Collider at Brookhaven National Lab and the Large Hadron Collider at CERN, the JET Collaboration has successfully extracted the jet transport coefficient of the QGP with high precision and carefully quantified uncertainties.

This shows how future studies of QGP properties can reach even higher levels of precision and sheds light on the mechanism causing the liquid-like nature of the plasma created in high-energy heavy-ion collisions.

More information: Karen M. Burke et al. Extracting the jet transport coefficient from jet quenching in high-energy heavy-ion collisions, *Physical Review C* (2014). [DOI: 10.1103/PhysRevC.90.014909](https://doi.org/10.1103/PhysRevC.90.014909)

Provided by US Department of Energy

Citation: Using fast particles to probe hot matter in nuclear collisions (2016, August 1) retrieved 27 April 2024 from <https://phys.org/news/2016-08-fast-particles-probe-hot-nuclear.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.