

Latest study of Tabby's star offers more weirdness

August 9 2016, by Bob Yirka



Allen Telescope Array. Credit: Seth Shostak, SETI Institute

A pair of researchers has added more evidence regarding the oddity of KIC 8462852, aka Tabby's Star. Benjamin Montet with the California Institute of Technology and Joshua Simon with Observatories of the

Carnegie Institution of Washington have uploaded a paper to the *arXiv* preprint server detailing their study of the star by analyzing data from the Kepler space observatory over the past four years. They found that the star has been decreasing in brightness at an unprecedented rate.

The tale of Tabby's Star began in September of last year when Louisiana State University astronomer Tabetha Boyajian reported anomalies in the unusual [light](#) curve of star KIC 8462852—over the years 2009 to 2013, its light appeared to dip in ways that did not conform to what would be expected if it were due to a planet passing in front of it, temporarily blocking some of its light. Her paper led to observations, commentaries and theories from others in the space community, though no one was able to come up with a reasonable explanation for what she had found. One researcher actually proposed that it might be due to alien activity. Then, earlier this year, Bradley Schaefer with Louisiana State University published results of his efforts studying photographic plates that had captured the star going back to the 19th century—he reported that the light from the star had dimmed 19 percent over just the past century. His report was not received warmly by all, as many suggested his data or approach was likely flawed.

But now, another team has found something similar. Montet and Simon studied images from the space-based Kepler observatory and found that light from Tabby's Star had decreased in brightness by approximately .34 percent a year for 1000 days starting in 2009, which was actually twice the rate that Schaefer had found. Even stranger, they found that over the next 200 days, the brightness of the star dimmed by another 2.5 percent before it finally leveled out.

Various scientists have offered possible explanations for the strange behavior of Tabby's star—from comet swarms to planetary remnants to the construction of a Dyson Sphere-like structure around the star to capture its energy by aliens—but so far, none of the theories has been

able to take into account all of the odd observations. That may change soon, however, as Boyajian, the astronomer who first noticed the star's strange behavior, ran a successful crowdsourcing campaign to pay for time at the Las Cumbres Observatory Global Telescope Network over the course of a year—this will give her a chance to catch the star in the act of blinking, allowing her to alert the rest of the astronomy community, which will presumably its sights on the star at once, and in so doing, perhaps solve the mystery.

More information: arxiv.org/abs/1608.01316

Abstract

KIC 8462852 is a superficially ordinary main sequence F star for which Kepler detected an unusual series of brief dimming events. We obtain accurate relative photometry of KIC 8462852 from the Kepler full frame images, finding that the brightness of KIC 8462852 monotonically decreased over the four years it was observed by Kepler. Over the first ~ 1000 days, KIC 8462852 faded approximately linearly at a rate of 0.341 ± 0.041 percent per year, for a total decline of 0.9%. KIC 8462852 then dimmed much more rapidly in the next ~ 200 days, with its flux dropping by more than 2%. For the final ~ 200 days of Kepler photometry the magnitude remained approximately constant, although the data are also consistent with the decline rate measured for the first 2.7 yr. Of a sample of 193 nearby comparison stars and 355 stars with similar stellar parameters, 0.6% change brightness at a rate as fast as 0.341 ± 0.041 percent per year, and none exhibit either the rapid decline by $>2\%$ or the cumulative fading by 3% of KIC 8462852. We examine whether the rapid decline could be caused by a cloud of transiting circumstellar material, finding while such a cloud could evade detection in sub-mm observations, the transit ingress and duration cannot be explained by a simple cloud model. Moreover, this model cannot account for the observed longer-term dimming. No known or proposed

stellar phenomena can fully explain all aspects of the observed light curve.

© 2016 Phys.org

Citation: Latest study of Tabby's star offers more weirdness (2016, August 9) retrieved 19 April 2024 from <https://phys.org/news/2016-08-latest-tabby-star-weirdness.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.