

A nanoscale wireless communication system via plasmonic antennas

August 25 2016



The pursuit of next-generation technologies places a premium on producing increased speed and efficiency with components built at scales small enough to function on a computer chip.

One of the barriers to advances in "on-chip" communications is the size of the electromagnetic waves at radio and microwave frequencies, which form the backbone of modern wireless technology. The relatively large waves handcuff further miniaturization.



Scientists trying to surpass these limitations are exploring the potential of optical conveyance that exploits the properties of much smaller wavelengths, such as those in the terahertz, infrared and visible frequencies.

A team of researchers at Boston College has developed the first nanoscale wireless communication system that operates at visible wavelengths using antennas that send and receive surface plasmons with an unprecedented degree of control, the team reports in the latest edition of the journal Nature's *Scientific Reports*.

Furthermore, the device affords an "in-plane" configuration, a prized class of two-way information transmission and recovery in a single path, according to the study, conducted by a team in the lab of Evelyn J. and Robert A. Ferris Professor of Physics Michael J. Naughton.

The findings mark an important first step toward a nanoscale version and visible frequency equivalent - of existing <u>wireless communication</u> <u>systems</u>, according to the researchers. Such on-chip systems could be used for high-speed communication, <u>high efficiency</u> plasmonic waveguiding and in-plane circuit switching - a process that is currently used in liquid crystal displays.

The device achieved communication across several wavelengths in tests using near-field scanning optical microscopy, according to lead coauthor Juan M. Merlo, a post-doctoral researcher who initiated the project.

"Juan was able to push it beyond the near field - at least to four times the width of a wavelength. That is true far-field transmission and nearly every device we use on a daily basis - from our cell phones to our cars relies on far-field transmission," said Naughton.



The device could speed the transmission of information by as much as 60 percent compared to earlier plasmonic waveguiding techniques and up to 50 percent faster than plasmonic nanowire waveguides, the team reports.

Surface plasmons are the oscillations of electrons coupled to the interface of an electromagnetic field and a metal. Among their unique abilities, surface plasmons can confine energy on that interface by fitting into spaces smaller than the waves themselves.

Researchers trying to exploit these subwavelength capabilities of surface plasmons have developed metallic structures, including plasmonic antennas. But a persistent problem has been the inability to achieve "in-line" containment of the emission and collection of the electromagnetic radiation.

The BC team developed a device with a three-step conversion process that changes a surface plasmon to a photon on transmission and then converts that elemental electromagnetic particle back to a surface plasmon as the receiver picks it up.

"We have developed a device where plasmonic antennas communicate with each other with photons transmitting between them," said Naughton. "This is done with high efficiency, with energy loss reduced by 50 percent between one antenna and the next, which is a significant enhancement over comparable architectures."

Central to the newfound control of the surface plasmons was the creation of a small gap of air between the waves and the silver surface of the device, said Merlo, who earned his PhD at Mexico's National Institute of Astrophysics, Optics and Electronics. By removing a portion of the glass substrate, the team reduced the disruptive pull of the material on the photons in transmission. Expanding and narrowing that gap proved



crucial to tuning the device.

With traditional silicon waveguides, dispersion reduces information transmission speed. Without that impediment, the new device capitalizes on the capability of <u>surface plasmons</u> to travel at 90 to 95 percent of the speed of light on a silver <u>surface</u> and photons traveling between the antennas at their inherent speed of light, Merlo said.

"Silicon-based optical technology has been around for years," said Merlo. "What we are doing is improving it to make it faster. We're developing a tool to make silicon photonics faster and greatly enhance rates of communication."

More information: Juan M. Merlo et al, Wireless communication system via nanoscale plasmonic antennas, *Scientific Reports* (2016). <u>DOI:</u> <u>10.1038/srep31710</u>

Provided by Boston College

Citation: A nanoscale wireless communication system via plasmonic antennas (2016, August 25) retrieved 2 May 2024 from <u>https://phys.org/news/2016-08-nanoscale-wireless-plasmonic-antennas.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.