


Scientists dig deep to cut mine poisons

August 5 2016



Associate Professor Sarah Harmer, centre, with postgraduate students Zoe Pettifer, Sian La Vars, Simon Lee, Belinda Bleeze and Connor Retallick at the Microbe Factory Laboratory at Flinders University

Widespread environmental damage caused by toxic chemicals and acid

leaching could be a thing of the past if new processing methods under development at Flinders University are scaled up and adopted by miners.

Scientists at Flinders in South Australia are developing a growing line-up of lab-based experiments to extract valuable minerals from ore using environmentally-friendly microbes, such as bacteria commonly found on mine locations.

"We're making real progress in finding better ways to more sustainably separate valuable ores such as copper, iron, lead and zinc," says ARC Future Fellow Associate Professor Sarah Harmer.

"We are using advanced technology to identify the specific naturally occurring bacterium which attach to each kind of mineral and modify its surface properties," Associate Professor Harmer says.

"This has the potential to be scaled up to one day replace toxic chemicals such as cyanide and xanthates now used to separate the minerals from ore in the early froth flotation phase of processing."

Waste rock and chemicals left behind from the process is usually saved in large tailing dams.

For the first time, Associate Professor Harmer's team at Flinders School of Chemical and Physical Sciences has used high-tech x-ray imaging and micro-spectroscopic methods at synchrotrons overseas to study the distribution of chemical species responsible for the selective attachment of bacteria and the separation of minerals.

They also have used the extreme light beams of the Australian Synchrotron and synchrotrons overseas to identify the chemical mechanisms of bioleaching of chalcopyrite, or copper ore, using soft and hard x-ray spectroscopies.

"The capability for microscale spectroscopic analysis of materials while in liquid under controlled electro-chemical conditions is helping to confirm which bacteria attach to specific mineral surfaces for the application of mineral leaching and separation of minerals without the use of toxic chemicals," Associate Professor Harmer explains.

"We have been testing the prototype electrochemical solution cell in collaboration with scientists from the Canadian Synchrotron, and we're looking forward to expanding that project in future."

Associate Professor Harmer says the small-scale experiments could be scaled up in trials by mining companies to extract ores more sustainably.

This would, in turn, start to reduce the toxicity of tailing dams on mine sites and long-term challenges of environmental remediation of mine sites around the world.

Toxic byproducts from mining and ore processing continue to plague the mining industry – and the environment.

Every year large volumes of heavy metals and [toxic chemicals](#) leach into soil, groundwater and rivers from mining company operations, as seen on Tasmania's West Coast to last year's devastating Samarco tailings dam collapse in Brazil.

Mining company Samarco, which is half-owned by BHP Billiton, has agreed to pay the Brazilian Government \$US2.3 billion (\$3.2 billion) for a dam collapse that killed 19 people in November last year.

Provided by Flinders University

Citation: Scientists dig deep to cut mine poisons (2016, August 5) retrieved 19 April 2024 from

<https://phys.org/news/2016-08-scientists-deep-poisons.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.