

# Scientists discover new function of FHY3 in plant meristem determinacy and maintenance

August 3 2016

---

In higher plants, meristems are responsible for the generation of all plant tissues and organs. While the shoot apical meristem (SAM) gives rise to all of the above-ground plant parts for the entire life of the plant, the floral meristem (FM) is terminated after the generation of all floral organs, a process known as FM determinacy, which helps ensure reproductive success, seed development, as well as the yield of agricultural crops.

A research Chinese research group has discovered a new function of far-red elongated hypocotyl 3 (FHY3) in plant meristem determinacy and maintenance by regulating sepallata 2 (SEP2) and clavata 3 (CLV3) expression.

It is well known that FHY3 plays pivotal roles in phyA signaling and circadian clock pathways as well as other developmental and physiological processes, including UV-B signaling, chloroplast biogenesis, chlorophyll biosynthesis, programmed cell death, ABA signaling, branching and flowering time at the plant vegetative stage. However, the roles of FHY3 in flower development remain unclear.

In this study, researchers isolated several *fhy3* mutations that enhanced the FM determinacy defects of *ag-10*, a weak agamous (*ag*) allele in a genetic screen. Genetic analysis showed that WUSCHEL (WUS) and CLV3, two central players in the establishment and maintenance of meristems, are epistatic to FHY3.

Using genome-wide ChIP-seq and RNA-seq data, researchers then identified hundreds of FHY3 target genes in flowers and found that FHY3 mainly acts as a transcriptional repressor in flower development, in contrast to its transcriptional activator role in seedlings. Binding motif-enrichment analyses indicated that FHY3 may co-regulate [flower development](#) with three flower-specific MADS-domain TFs and four basic helix-loop-helix TFs that are involved in photomorphogenesis.

Further study revealed that CLV3, sepallata 1 (SEP1), and SEP2 are FHY3 target genes. In SAM, FHY3 directly represses CLV3, which consequently regulates WUS to maintain the stem cell pool. Intriguingly, CLV3 expression did not change significantly in FHY3 and phytochrome B mutants before and after light treatment, indicating that FHY3 and phytochrome B are involved in light-regulated meristem activity. In FM, FHY3 directly represses CLV3, but activates SEP2, to ultimately promote FM determinacy.

This work reveals new insights into the mechanisms of meristem maintenance and determinacy, and illustrates how the roles of a single TF may vary between organs and developmental stages. More interestingly, FHY3 may act as a bridge molecule in the cross-talk between external signals and endogenous cues to coordinate plant development.

The article has been published in *Proceedings of the National Academy of Sciences*.

**More information:** Dongming Li et al, activatesbut inhibitsto regulate meristem determinacy and maintenance in, *Proceedings of the National Academy of Sciences* (2016). [DOI: 10.1073/pnas.1602960113](https://doi.org/10.1073/pnas.1602960113)

Provided by Chinese Academy of Sciences

Citation: Scientists discover new function of FHY3 in plant meristem determinacy and maintenance (2016, August 3) retrieved 30 June 2024 from

<https://phys.org/news/2016-08-scientists-function-fhy3-meristem-determinacy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.