

Scientists describe a new transmission path of a tropical disease

August 2 2016



Two flies sucking lesion exudate on a skin ulcer from a Treponema pallidum infected olive baboon (Papio anubis) at Lake Manyara National Park in Tanzania. Credit: F. Paciencia

Lesions on arms and legs, deformed faces – yaws is a tropical disease that infects the skin, bones and cartilage. It is caused by the bacterium



Treponema pallidum subspecies pertenue. Mostly children in remote tropical areas of Africa, Southeast Asia and the Pacific are infected. Until recently it was assumed that the disease is only spread by direct skin contact with an infected person. An international research group led by Sascha Knauf from the German Primate Center – Leibniz Institute for Primate Research now published a study that suggests flies as mechanical vectors for the bacterium. The finding is of importance for currently ongoing yaws eradication campaigns.

Where the roads end and only scattered villages interrupt the jungle, the area of yaws begins. In its primary stage, yaws infection causes skin lesions and if untreated progresses into severe bone and joint deformations of the extremities and face. Infection is transmitted from person-to-person by direct contact with contagious lesions. The World Health Organization (WHO) assumes that 75 to 80 per cent of the patients are under 15 years old.

Theoretically, one pill of an antibiotic would be enough to prevent children from suffering and to cure the disease that is caused by the bacterium Treponema pallidum subspecies pertenue. In practice however, there are many villages that do not have access to modern medicine. The WHO, for the second time, tries to eradicate yaws finally by 2020. However, as studies by Sascha Knauf and colleagues suggest, there could be major challenges involved. The scientists have not only isolated an identical bacterium in monkeys suggesting that there is a nonhuman reservoir, they now also discovered the pathogen on <u>flies</u>. "The human-to-human as well as possible monkey-to-human transmission becomes a complete new dimension," says Sascha Knauf from the German Primate Center.

Sascha Knauf and colleagues investigated 207 flies originating from two national parks in Tanzania. In both parks skin ulcerations caused by Treponema are common in wild baboons. In about 20 per cent of the



wild-caught flies Treponema-DNA was found. "Our results support the possibility that flies play a role in yaws transmission", says Sascha Knauf, first author of the publication. The scientists want to continue their research on the topic, because it is not yet clear, whether the bacteria found in the flies are viable. "If we can confirm that the flies transmit viable bacteria and that infected monkeys act as a nonhuman reservoir, there will always be cases of yaws even if the disease is apparently eradicated due to medicine and hygienic measures," says Knauf.





A male olive baboon infected with Treponema pallidum. Credit: Sascha Knau

More information: Sascha Knauf et al. Isolation of Treponema DNA from Necrophagous Flies in a Natural Ecosystem, *EBioMedicine* (2016). DOI: 10.1016/j.ebiom.2016.07.033



Provided by The German Primate Center

Citation: Scientists describe a new transmission path of a tropical disease (2016, August 2) retrieved 26 April 2024 from https://phys.org/news/2016-08-scientists-transmission-path-tropical-disease.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.