

Toxic blue-green algae adapt to rising CO₂

August 4 2016



Credit: Universiteit van Amsterdam (UVA)

A common type of blue-green algae is finding it easy to adapt to Earth's rising CO₂ levels, meaning blue-green algae – of which there are many toxin-producing varieties – are even more adept at handling changing

climatic conditions than scientists previously supposed. A team of microbiologists at the University of Amsterdam (UvA) are reporting this finding in the journal *PNAS* this week, and point here at implications for clean drinking water, swimming safety and freshwater ecosystems.

The research team, led by Professor of Aquatic Microbiology Jef Huisman, trained their microscopes on *Microcystis*, a type of blue-green algae that proliferate in lakes and reservoirs in summer. The team analysed the genetic composition of [cyanobacteria](#) (blue-green algae's scientific name), observing *Microcystis* in both the lab and the Kennemer lake, under CO₂-rich and poor conditions. "Before this, the adaptive potential of these harmful cyanobacteria in response to increasing CO₂ concentrations had never been studied systematically, even though this can help us predict how algal blooms will develop in future", explains Xing Ji, a PhD researcher on the team.

In both the lab and the lake, cyanobacteria's genetic makeup changed in response to increasing CO₂ concentrations. "It's a textbook example of natural selection", says lead author Giovanni Sandrini. "Cyanobacteria absorb CO₂ during photosynthesis to produce their biomass, and we observed that the strain best equipped to absorb dissolved CO₂ eventually gains the upper hand."

Some *Microcystis* strains have a slow but efficient carbon uptake system that enables them to squeeze out the last bit of CO₂ from the water even at very low concentrations. Those strains become dominant in low CO₂ conditions. By contrast, other strains have a fast uptake system that allows them to take up dissolved CO₂ at very high rates when in high concentrations. "We discovered that these high-speed strains enjoy a major selective advantage in CO₂-rich water", Sandrini continues. "Given the rising atmospheric CO₂ values, these strains are poised to thrive."

Bathing and drinking water

Cyanobacteria's adaptation to rising CO₂ is cause for concern. That's because *Microcystis* can produce microcystin, a toxin that causes liver damage in birds and mammals. In high concentrations, cyanobacteria also disrupt [freshwater ecosystems](#), killing fish and aquatic plants. In the Netherlands, blue-green algal blooms regularly put swimming areas off limits.

Ji personally experienced just how harmful these bacteria have already proved to be in 2007, when he was living in eastern China, where cyanobacteria covered the entire surface of Lake Taihu, a 2000-km² lake, and led to a [drinking water](#) crisis affecting five million people. "I watched my mother arguing with other supermarket shoppers who all had their sights set on the last bottles of drinking water. It's precisely because I'm aware of how poor water quality can impact society that I am happy to be doing research that can yield relevant insights."

More information: Giovanni Sandrini et al. Rapid adaptation of harmful cyanobacteria to rising CO₂, *Proceedings of the National Academy of Sciences* (2016). [DOI: 10.1073/pnas.1602435113](https://doi.org/10.1073/pnas.1602435113)

Provided by University of Amsterdam

Citation: Toxic blue-green algae adapt to rising CO₂ (2016, August 4) retrieved 29 June 2024 from <https://phys.org/news/2016-08-toxic-blue-green-algae-co2.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.