


# One billion stars in 3-D: Gaia's billion-star map hints at treasures to come

September 14 2016

---



An all-sky view of stars in our Galaxy – the Milky Way – and neighbouring galaxies, based on the first year of observations from ESA's Gaia satellite, from July 2014 to September 2015. Credit: ESA/Gaia/DPAC

The first catalogue of more than a billion stars from ESA's Gaia satellite was published today – the largest all-sky survey of celestial objects to

date.

On its way to assembling the most detailed 3-D map ever made of our Milky Way galaxy, Gaia has pinned down the precise position on the sky and the brightness of 1142 million stars.

As a taster of the richer catalogue to come in the near future, today's release also features the distances and the motions across the sky for more than two million stars.

"Gaia is at the forefront of astrometry, charting the sky at precisions that have never been achieved before," says Alvaro Giménez, ESA's Director of Science.

"Today's release gives us a first impression of the extraordinary data that await us and that will revolutionise our understanding of how stars are distributed and move across our Galaxy."

Launched 1000 days ago, Gaia started its scientific work in July 2014. This first release is based on data collected during its first 14 months of scanning the sky, up to September 2015.

"The beautiful map we are publishing today shows the density of stars measured by Gaia across the entire sky, and confirms that it collected superb data during its first year of operations," says Timo Prusti, Gaia project scientist at ESA.



Artist's impression of Gaia mapping the stars of the Milky Way. Credit: ESA/ATG medialab; background: ESO/S. Brunier

The stripes and other artefacts in the image reflect how Gaia scans the sky, and will gradually fade as more scans are made during the five-year mission.

"The satellite is working well and we have demonstrated that it is possible to handle the analysis of a billion stars. Although the current data are preliminary, we wanted to make them available for the astronomical community to use as soon as possible," adds Dr Prusti.

Transforming the raw information into useful and reliable stellar positions to a level of accuracy never possible before is an extremely

complex procedure, entrusted to a pan-European collaboration of about 450 scientists and software engineers: the Gaia Data Processing and Analysis Consortium, or DPAC.

"Today's release is the result of a painstaking collaborative work over the past decade," says Anthony Brown from Leiden University in the Netherlands, and consortium chair.

"Together with experts from a variety of disciplines, we had to prepare ourselves even before the start of observations, then treated the data, packaged them into meaningful astronomical products, and validated their scientific content."

In addition to processing the full billion-star catalogue, the scientists looked in detail at the roughly two million stars in common between Gaia's first year and the earlier Hipparcos and Tycho-2 Catalogues, both derived from ESA's Hipparcos mission, which charted the sky more than two decades ago.

By combining Gaia data with information from these less precise catalogues, it was possible to start disentangling the effects of 'parallax' and 'proper motion' even from the first year of observations only. Parallax is a small motion in the apparent position of a star caused by Earth's yearly revolution around the Sun and depends on a star's distance from us, while proper motion is due to the physical movement of stars through the Galaxy.

In this way, the scientists were able to estimate distances and motions for the two million stars spread across the sky in the combined Tycho–Gaia Astrometric Solution, or TGAS.

This new catalogue is twice as precise and contains almost 20 times as many stars as the previous definitive reference for astrometry, the



An all-sky view of stars in our Galaxy – the Milky Way – and neighbouring galaxies, based on the first year of observations from ESA's Gaia satellite, from July 2014 to September 2015. Credit: ESA/Gaia/DPAC

For the closest 14 open clusters, the new data reveal many stars surprisingly far from the centre of the parent cluster, likely escaping to populate other regions of the Galaxy."

Many more stellar clusters will be discovered and analysed in even greater detail with the extraordinary data that Gaia continues to collect and that will be released in the coming years.

The new stellar census also contains 3194 [variable stars](#), stars that rhythmically swell and shrink in size, leading to periodic brightness changes.

Many of the variables seen by Gaia are in the Large Magellanic Cloud, one of our galactic neighbours, a region that was scanned repeatedly during the first month of observations, allowing accurate measurement of their changing brightness.

Details about the brightness variations of these stars, 386 of which are new discoveries, are published as part of today's release, along with a first study to test the potential of the data.

"Variable stars like Cepheids and RR Lyraes are valuable indicators of cosmic distances," explains Gisella Clementini from INAF and the Astronomical Observatory of Bologna, Italy.

"While parallax is used to measure distances to large samples of stars in



the Milky Way directly, variable stars provide an indirect, but crucial step on our 'cosmic distance ladder', allowing us to extend it to faraway galaxies."

This is possible because some kinds of variable stars are special. For example, in the case of Cepheid stars, the brighter they are intrinsically, the slower their brightness variations. The same is true for RR Lyraes when observed in infrared light. The variability pattern is easy to measure and can be combined with the apparent brightness of a star to infer its true brightness.

This is where Gaia steps in: in the future, scientists will be able to determine very accurate distances to a large sample of variable stars via Gaia's measurements of parallaxes. With those, they will calibrate and improve the relation between the period and brightness of these stars, and apply it to measure distances beyond our Galaxy. A preliminary application of data from the TGAS looks very promising.


"This is only the beginning: we measured the distance to the Large Magellanic Cloud to test the quality of the data, and we got a sneak preview of the dramatic improvements that Gaia will soon bring to our understanding of cosmic distances," adds Dr Clementini.

Knowing the positions and motions of stars in the sky to astonishing precision is a fundamental part of studying the properties and past history of the Milky Way and to measure distances to stars and galaxies, but also has a variety of applications closer to home – for example, in the Solar System.

In July, Pluto passed in front of a distant, faint star, offering a rare chance to study the atmosphere of the dwarf planet as the star gradually disappeared and then reappeared behind Pluto.







On 19 July 2016, Pluto passed in front of the faint star UCAC4 345-180315, offering a rare chance to study the atmosphere of the dwarf planet as the star first gradually disappeared and then reappeared behind Pluto. Credit: B. Sicardy (LESIA, Observatoire de Paris, France), P. Tanga (Observatoire de la Côte d'Azur, Nice, France), A. Carbognani (Osservatorio Astronomico Valle d'Aosta, Italy), Rodrigo Leiva (LESIA, Observatoire de Paris)

This stellar occultation was visible only from a narrow strip stretching across Europe, similar to the totality path that a solar eclipse lays down on our planet's surface. Precise knowledge of the star's position was crucial to point telescopes on Earth, so the exceptional early release of the Gaia position for this star, which was 10 times more precise than previously available, was instrumental to the successful monitoring of this rare event.

Early results hint at a pause in the puzzling pressure rise of Pluto's tenuous atmosphere, something that has been recorded since 1988 in spite of the dwarf planet moving away from the Sun, which would suggest a drop in pressure due to cooling of the atmosphere.

"These three examples demonstrate how Gaia's present and future data will revolutionise all areas of astronomy, allowing us to investigate our place in the Universe, from our local neighbourhood, the Solar System, to Galactic and even grander, cosmological scales," explains Dr Brown.

This first data release shows that the mission is on track to achieve its ultimate goal: charting the positions, distances, and motions of one billion [stars](#) – about 1% of the Milky Way's stellar content – in three dimensions to unprecedented accuracy.

"The road to today has not been without obstacles: Gaia encountered a number of technical challenges and it has taken an extensive collaborative effort to learn how to deal with them," says Fred Jansen, Gaia mission manager at ESA.

"But now, 1000 days after launch and thanks to the great work of everyone involved, we are thrilled to present this first dataset and are looking forward to the next release, which will unleash Gaia's potential to explore our Galaxy as we've never seen it before."

**More information:** The data from Gaia's first release can be accessed at [archives.esac.esa.int/gaia](http://archives.esac.esa.int/gaia)

Fifteen scientific papers describing the data contained in the release and their validation process will appear in a special issue of [Astronomy & Astrophysics](#).

Provided by European Space Agency

Citation: One billion stars in 3-D: Gaia's billion-star map hints at treasures to come (2016, September 14) retrieved 13 March 2024 from <https://phys.org/news/2016-09-billion-stars-d-gaia-billion-star.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--