

# **Gecko study offers evidence that small morphological changes can lead to large changes in function**

September 29 2016, by Iqbal Pittalwala

---



G. humeralis in Trinidad. Credit: Timothy Higham, UC Riverside

How do key innovations in the animal kingdom arise? To explore this question, gecko expert Timothy Higham, an associate professor of biology at the University of California, Riverside, led a team of evolutionary biologists to study *Gonatodes*, a genus of dwarf geckos. In the process, the researchers found a gecko, *Gonatodes humeralis*, that they posit offers a "snapshot" into the evolution of adhesion in geckos.

"The gecko adhesive apparatus, one of the most spectacular innovations displayed by vertebrates, has been intensively studied for the last 16 years and is of considerable interest to nanotechnologists and [biomimeticists](#)," Higham said. "But almost nothing is known about the origin of this adhesive capability. *G. humeralis*, found in South America, shows how the adhesive capabilities of [geckos](#) may have come about. Our integrative analysis of this gecko shows that unexpectedly it has microscopic hairs, called setae, underneath its toes, which allow it to do something dramatically different than all other geckos in the *Gonatodes* genus: cling to smooth surfaces such as leaves. It does this without all of the complex structure of the toes that typify the geckos that we are more familiar with. In the lab, this [gecko can climb smooth vertical surfaces](#) using its incipient adhesive system."

Higham explained that the setae interact with surfaces through attractive van der Waals forces. The relatively simple expression of setae on the digits of *G. humeralis* thus provide an enormous advantage in sectors of the habitat typified by smooth, low-friction, inclined surfaces, such as leaves and slippery stems, allowing *G. humeralis* to avoid predators by occupying habitat that other members of the genus cannot. While it can

securely attach to vertical bamboo shoots, for example, other species in the *Gonatodes* genus generally scale rough tree trunks, rocks, fallen palm trees and move on the ground – areas where their predators abound.

"The relatively simple adhesive system of the *G. humeralis* is indicative that slight modifications in form can dramatically influence functional outcomes and the ecological niches that can be exploited," Higham said.

"This ostensibly padless gecko offers us a snapshot – a crucial intermediate stage – of the evolution of the adhesion apparatus. It's telling us, 'Look, this is how pad-bearing geckos started to acquire adhesion.'"

Further, the findings indicate that the origin of adhesion in geckos was gradual and led to major shifts in ecology and function. They suggest, too, that subtle morphological changes are able to trigger rapid evolution.



G. humeralis in French Guiana. Credit: Timothy Higham, UC Riverside

Study results appear Sept. 29 in the *Biological Journal of the Linnean Society*.

The setae of *G. humeralis* are short and simple compared to those of pad-bearing geckos, such as [tokay geckos](#). The setae are located adjacent to friction-enhancing "spinules" – small projections, which play no role in adhesion, that are found underneath the feet of many lizards and geckos. The authors argue that the setae of *G. humeralis* result from a transformation of the spinules.

"Until now, we had not seen a gecko showing the beginnings of the adhesive system," Higham said. "In all the innovations seen in the [animal kingdom](#), we rarely get to see their beginnings. Our findings serve as good evidence against intelligent design ideas. Evolution takes place in incremental steps, as the 'snapshot' we report on shows. Complexity does not start with complexity. Small modifications can, however, lead to complexity. Key innovations can come about in small incremental steps and lead to feedback processes that result in the more complex renditions of such systems. Our research offers more experimental evidence to show this is true."

Higham and his colleagues found *G. humeralis* specimens in Trinidad and French Guiana. They first used scanning electron microscopy to examine the microanatomy underneath the geckos' toes and morphometrics to compare them to close relatives. After these initial observations, the team studied this species and others in the laboratory using a combination of high-speed video to measure locomotion and force transducers to quantify clinging ability, finding that only *G. humeralis* could generate clinging force and climb vertical smooth

acrylic.

"Our work would interest researchers in nanotechnology and biomimicry because it explores the question: 'What are the minimum structural modifications to enable adhesion?' Gecko-inspired robots tend to have fully developed adhesion systems. But is that necessary? Our work suggests that not much is needed to get a good adhesion system going, and may therefore help simplify how we approach biomimicry and how we recreate [adhesion](#) in the lab."

**More information:** Timothy E. Higham et al. On the origin of frictional adhesion in geckos: small morphological changes lead to a major biomechanical transition in the genus, *Biological Journal of the Linnean Society* (2016). [DOI: 10.1111/bij.12897](https://doi.org/10.1111/bij.12897)

Provided by University of California - Riverside

Citation: Gecko study offers evidence that small morphological changes can lead to large changes in function (2016, September 29) retrieved 1 May 2024 from <https://phys.org/news/2016-09-gecko-evidence-small-morphological-large.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--