


# **Novel study identifies an area of the mosquito brain that mixes taste and smell**

October 10 2016

---



A female *Anopheles gambiae* mosquito with olfactory neurons on the antennae, maxillary palp and labella labeled in green. Credit: Olena Riabinina and Courtney Akitake, Johns Hopkins Medicine

A new study by Johns Hopkins researchers suggests that a specialized area of the mosquito brain mixes tastes with smells to create unique and preferred flavors. The findings advance the possibility, they say, of identifying a substance that makes "human flavor" repulsive to the malaria-bearing species of the mosquitoes, so instead of feasting on us, they keep the disease to themselves, potentially saving an estimated 450,000 lives a year worldwide.

A report on the research appeared online on Oct. 3 in the journal *Nature Communications*. Malaria is an infectious parasite disease of humans and animals transmitted by the bite of the female *Anopheles gambiae* mosquito. In 2015, experts estimate it affected 214 million people, mostly in Africa, despite decades of mosquito eradication and control efforts. There is no malaria vaccine, and although the disease is curable in early stages, treatment is costly and difficult to deliver in places where it is endemic.

"All mosquitoes, including the one that transmits malaria, use their sense of smell to find a host for a blood meal. Our goal is to let the mosquitoes tell us what smells they find repulsive and use those to keep them from biting us," says Christopher Potter, Ph.D., assistant professor of neuroscience at the Johns Hopkins University School of Medicine.

Because smell is essential to mosquito survival, each mosquito has three pairs of "noses" for sensing odors: two antennae, two maxillary palps and

two labella. The maxillary palps are thick, fuzzy appendages that protrude from the lower region of the mosquito's head, more or less parallel to its proboscis, the long, flexible sheath that keeps its "feeding needle" under wraps until needed. At the very tip of the proboscis are the labella, two small regions that contain both "gustatory" neurons that pick up tastes and [olfactory neurons](#) for recognizing odorants.

To better understand how *An. gambiae* mosquitoes that cause malaria receive and process olfactory information from so many sensory regions, Potter's team wanted to see where olfactory neurons from those regions go to in the brain.

They used a powerful genetic technique—never before accomplished in mosquitoes, according to Potter—to make certain neurons "glow" green. The green glowing label was designed to appear specifically in neurons that receive complex odors through proteins called odorant receptors (ORs), since OR neurons are known to help distinguish humans from other warm-blooded animals in *Aedes aegypti* mosquitoes, which carry the Zika virus.

"This is the first time researchers managed to specifically target [sensory neurons](#) in mosquitoes. Previously, we had to use flies as a proxy for all insects, but now we can directly study the sense of smell in the insects that spread malaria," says Olena Riabinina, Ph.D., the lead author of the study and a postdoctoral fellow now at the Imperial College London. "We were pleasantly surprised by how well our genetic technique worked and how easy it is now to see the smell-detecting neurons. The ease of identification will definitely simplify our task of studying these neurons in the future."

As expected, Potter says, the OR neurons from the antennae and maxillary palps went to symmetrical areas of the brain called antennal lobes, just as they do in flies. But Potter was quite surprised when he saw

that the OR neurons from the labella went to the so-called subesophageal zone, which, he says, had never before been associated with the sense of smell in any fly or insect; it had only been associated with the sense of taste.

"That finding suggests that perhaps mosquitoes don't just like our smell, but also our flavor," says Potter. "It's likely that the odorants coming off our skin are picked up by the labella and influence the preferred taste of our skin, especially when the mosquito is looking for a place to bite."

Potter says the finding potentially offers researchers one more way to repel mosquitoes. The antennae and maxillary palps are more specialized for picking up long-range signals, while the labella come into direct contact with our skin. In fact, Potter says, before injecting their needlelike proboscis, mosquitoes use the labella to probe about on a victim's skin. "We don't really know why they do that, but we suspect that they're looking for sensory cues that hint at easy access to a blood vessel," he says. "This suggests that a combination of repellants could keep mosquitoes from biting us in two ways. One could target the antennal neurons and reduce the likelihood that they come too close, while another could target the labellar neurons and make the mosquitoes turn away in disgust—before sucking our blood—if they got close enough to land on us."

The two-part genetic system Potter devised to generate the glowing neurons will make it much easier for his and other laboratories to mix and match [genetically altered mosquitoes](#) to generate new traits, he says. His group has already created a strain of *An. gambiae* mosquitoes whose OR neurons glow green upon activation. Scientists can thus see which neurons light up in response to a specific smell.

"Using this method, we hope to find an odorant that is safe and pleasant-smelling for us but strongly repellant to mosquitoes at very low

concentrations," says Potter.

His group was also able to compare the brains of male and female mosquitoes. Since only females use their sense of smell to find humans and males feed only on nectar, it was previously thought that males had just a rudimentary sense of smell. The Potter group found instead that males have the same level of complexity in their brains to detect odors as females but have fewer olfactory neurons. "It appears that males might just have a scaled-down version of a female's sense of smell. So they can still smell everything a female smells, just not as well," Potter says.

His group plans to study other types of neurons to better understand how signals from the [mosquitoes'](#) three types of olfactory receptors interact to influence their behavior. For example, why is lactic acid not attractive on its own but highly attractive when mixed with carbon dioxide?

"We'd like to figure out what regions and [neurons](#) in the brain lead to this combined effect," says Potter. "If we can identify them, perhaps we could also stop them from working."

**More information:** Olena Riabinina et al, Organization of olfactory centres in the malaria mosquito *Anopheles gambiae*, *Nature Communications* (2016). [DOI: 10.1038/NCOMMS13010](https://doi.org/10.1038/NCOMMS13010)

Provided by Johns Hopkins University School of Medicine

Citation: Novel study identifies an area of the mosquito brain that mixes taste and smell (2016, October 10) retrieved 16 August 2024 from <https://phys.org/news/2016-10-area-mosquito-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private
---

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.