

'Weighing' atoms with electrons

October 11 2016



Although atoms in graphene can be "seen" by shooting electrons through the material in what is called transmission electron microscopy, different isotopes can appear identical. Credit: Copyright: Koponen+Hildén, Creative Commons BY 4.0

The different elements found in nature each have their distinct isotopes. For carbon, there are 99 atoms of the lighter stable carbon isotope 12C for each 13C atom, which has one more neutron in its nucleus. Apart from this natural variation, materials can be grown from isotope-



enriched chemicals. This allows scientists to study how the atoms arrange into solids, for example to improve their synthesis. Yet, most traditional techniques to measure the isotope ratio require the decomposition of the material or are limited to a resolution of hundreds of nanometers, obscuring important details.

In the new study, led by Jani Kotakoski, the University of Vienna researchers used the advanced scanning transmission electron microscope Nion UltraSTEM100 to measure isotopes in nanometersized areas of a graphene sample. The same energetic electrons that form an image of the graphene structure can also eject one atom at a time due to scattering at a carbon nucleus. Because of the greater mass of the 13C isotope, an electron can give a 12C atom a slightly harder kick, knocking it out more easily. How many electrons are on average required gives an estimate of the local isotope concentration. "The key to making this work was combining accurate experiments with an improved theoretical model of the process", says Toma Susi, the lead author of the study.

Publishing in *Nature Communications* allowed the team to fully embrace open science. In addition to releasing the peer review reports alongside the article, a comprehensive description of the methods and analyses is included. However, the researchers went one step further and uploaded their microscopy data onto the open repository figshare. Anyone with an Internet connection can thus freely access, use and cite the gigabytes of high-quality images. Toma Susi continues: "To our knowledge, this is the first time electron microscopy data have been openly shared at this scale."

The results show that atomic-resolution electron microscopes can distinguish between different isotopes of carbon. Although the method was now demonstrated only for graphene, it can in principle be extended for other two-dimensional materials, and the researchers have a patent pending on this invention. "Modern microscopes already allow us to



resolve all atomic distances in solids and to see which chemical elements compose them. Now we can add isotopes to the list", Jani Kotakoski concludes.



The lighter the atom, the fewer electrons are on everage needed to eject it. Credit: Copyright: Koponen+Hildén, Creative Commons BY 4.0

More information: Toma Susi et al, Isotope analysis in the transmission electron microscope, *Nature Communications* (2016). <u>DOI:</u> <u>10.1038/ncomms13040</u>

Open data: Atomic resolution electron irradiation time series of isotopically labeled monolayer graphene: Toma Susi, Christoph Hofer, Giacomo Argentero, Gregor T. Leuthner, Timothy J. Pennycook,



Clemens Mangler, Jannik C. Meyer & Jani Kotakoski. figshare (2016). DOI: 10.6084/m9.figshare.c.3311946.v1

Provided by University of Vienna

Citation: 'Weighing' atoms with electrons (2016, October 11) retrieved 24 May 2024 from https://phys.org/news/2016-10-atoms-electrons.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.