


# Diamonds aren't forever: Team create first quantum computer bridge

October 14 2016

---



An array of holes (purple) etched in diamond, with two silicon atoms (yellow) placed between the holes. Credit: Sandia National Laboratories

By forcefully embedding two silicon atoms in a diamond matrix, Sandia researchers have demonstrated for the first time on a single chip all the

components needed to create a quantum bridge to link "People have already built small quantum computers," says Sandia researcher Ryan Camacho. "Maybe the first useful one won't be a single giant quantum computer but a connected cluster of small ones."

Distributing quantum information on a bridge, or network, could also enable novel forms of quantum sensing, since quantum correlations allow all the atoms in the network to behave as though they were one single atom.

The joint work with Harvard University used a focused [ion beam](#) implanter at Sandia's Ion Beam Laboratory designed for blasting single ions into precise locations on a diamond substrate. Sandia researchers Ed Bielejec, Jose Pacheco and Daniel Perry used implantation to replace one carbon atom of the diamond with the larger silicon atom, which causes the two [carbon atoms](#) on either side of the silicon atom to feel crowded enough to flee. That leaves the silicon atom a kind of large landowner, buffered against stray electrical currents by the neighboring non-conducting vacancies.

Though the silicon atoms are embedded in a solid, they behave as though floating in a gas, and therefore their electrons' response to quantum stimuli are not clouded by unwanted interactions with other matter.

"What we've done is implant the silicon atoms exactly where we want them," said Camacho. "We can create thousands of implanted locations, which all yield working quantum devices, because we plant the atoms well below the surface of the substrate and anneal them in place. Before this, researchers had to search for emitter atoms among about 1,000 randomly occurring defects—that is, non-carbon atoms—in a diamond substrate of a few microns to find even one that emitted strongly enough to be useful at the single photon level."

Once the [silicon atoms](#) are settled in the diamond substrate, laser-generated photons bump silicon electrons into their next higher atomic energy state; when the electrons return to the lower energy state, because all things seek the lowest possible energy level, they spit out quantized photons that carry information through their frequency, intensity and the polarization of their wave.

"Harvard researchers performed that experiment, as well as the optical and quantum measurements," said Camacho. "We did the novel device fabrication and came up with a clever way to count exactly how many ions are implanted into the diamond substrate."

Sandia researcher John Abraham and other Sandia researchers developed special detectors—metal films atop the diamond substrate—that showed the ion beam implants were successful by measuring the ionization signal produced by single ions.

"Pretty cool, huh?" said Camacho.

The journal *Science* thought so. The work is published in the current issue.

**More information:** A. Sipahigil et al, An integrated diamond nanophotonics platform for quantum optical networks, *Science* (2016). [DOI: 10.1126/science.aah6875](https://doi.org/10.1126/science.aah6875)

Provided by Sandia National Laboratories

Citation: Diamonds aren't forever: Team create first quantum computer bridge (2016, October 14) retrieved 25 April 2024 from <https://phys.org/news/2016-10-diamonds-team-quantum-bridge.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.