


Gamma ray camera offers new view on ultra-high energy electrons in plasma

October 27 2016



Cross section of gamma ray imager and illustration of emission from energetic electrons. Credit: General Atomics


Researchers at General Atomics (GA) have invented a new kind of gamma ray camera that can image beams of energetic electrons inside ultra-hot fusion plasma.

The device is used in ongoing global research that is developing fusion into a new clean energy source. Turning fusion fuel into extractable

energy requires it to be hotter than the center of the sun, hence in the plasma state. If the shutdown phase of operation is not controlled well, released magnetic energy can drive a population of electrons to relativistic speeds. If this population is not controlled, the electrons impact the inner walls of the plasma chamber, leading to material damage.

A team of researchers is working to better understand the properties of these electrons at the DIII-D National Fusion Facility operated by GA in San Diego for the U.S. Department of Energy. They designed and built a Gamma Ray Imager to capture the image of these particles.

The Gamma Ray Imager works on the principle of a standard pinhole camera (Figure 1), except that it is made of lead and weighs 420 pounds (190.5 kilograms). The imager actually records images of equally energetic [gamma rays](#) that are emitted by the electrons, and the lead is necessary to achieve a good focus (Figure 2). These gamma ray measurements provide information about the energy, direction, and quantity of electrons in the relativistic population, giving researchers an unparalleled view of how the [energetic electrons](#) evolve and interact with the fusion plasma.




Each pixel of the camera is an individual detector pictured above. These are arranged to look into the plasma and focus on the hottest part of the plasma.
Credit: General Atomics

"This system allows us to see with unprecedented detail how different plasma properties can mitigate these electrons," said Dr. Carlos Paz-Soldan, the scientist who led the first experiments utilizing the new camera. The results, to be presented at the American Physical Society Division of Plasma Physics conference Oct. 31-Nov. 4, demonstrate experimentally that radiation "reaction" forces are able to sap the highest energy electrons while collisions with other electrons are most effective at low energy (Figure 3).

These measurements imply that energetic electron control is not one-size-fits all, and that different energies require different control strategies.

With the new measurements, scientists can compare the behavior of the electron populations to theoretical models being developed by research

teams worldwide. These models are, in turn, crucial to predict how the electron populations will behave in new reactors, such as the ITER tokamak now under construction in Cadarache, France, and thus ensure they can be adequately controlled.



Electron energy distribution changes as the radiation back-reaction force is increased by raising the magnetic field. Credit: General Atomics

More information: meetings.aps.org/Meeting/DPP16/Session/CO4.10

Provided by American Physical Society

Citation: Gamma ray camera offers new view on ultra-high energy electrons in plasma (2016, October 27) retrieved 8 September 2024 from <https://phys.org/news/2016-10-gamma-ray-camera-view-ultra-high.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.