

Ultrastructure of a condensed chromosome-like structure in a cyanobacterium

October 21 2016

Eukaryotic cells, including human cells, form paired condensed chromosomes before cell division. The paired chromosomes are then equally divided into daughter cells. Prokaryotic cells, including bacteria, do not have such a DNA distribution system.

Researchers working at the National Institute for Physiological Sciences (NIPS) and Saitama University have discovered that the photosynthetic cyanobacterium *Synechococcus elongatus* shows eukaryotic condensed chromosome-like DNA compaction prior to [cell division](#), and were able to reveal details of the transiently formed structure. The bacteria divide at the moment of transition from day to night if cultivated under rigorous 12-hour light/12-hour dark cycles. The ephemeral 3-D structure of the compact DNA was captured by high-voltage cryo-electron tomography following [rapid freezing](#). The researchers show that the structure is similar to the condensed chromosomes in [eukaryotic cells](#). The compacted DNA also includes many small, paired polyphosphate bodies, some of which seem to maintain contact with DNA that appears to twist away from them, indicating that they may act as suppliers and regulators of phosphate for DNA synthesis. These new findings may deepen our understanding of how the eukaryotic cell arrived at a system using condensed chromosomes in the long evolution of biological cells.

For the study, published in *Scientific Reports*, the researchers used a combination of rigorous synchronized cultivation and sophisticated imaging by high-voltage cryo-electron tomography after rapid freezing to depict the transient 3-D structure of the compacted DNA in

cyanobacteria as thick as 1 μm . The polyphosphate bodies included in the compacted DNA may play a role as phosphate suppliers for DNA synthesis, which are duplicated in tandem with DNA.

"We know that the difference between prokaryotes and eukaryotes is in the DNA packaging and delivering system, where eukaryotes have developed a nuclear membrane to protect their genes and form condensed chromosomes to properly parcel the genome into the daughter cell, whereas prokaryotes do not have such a system," says corresponding author Kazuyoshi Murata. "Our new findings throw light on the duplication and segregation mechanisms of cyanobacterial DNA and point to an important role for polyphosphate bodies. They may give hints as to by what steps the evolution from prokaryote to eukaryote took place."

More information: Kazuyoshi Murata et al, Ultrastructure of compacted DNA in cyanobacteria by high-voltage cryo-electron tomography, *Scientific Reports* (2016). [DOI: 10.1038/srep34934](https://doi.org/10.1038/srep34934)

Provided by National Institutes of Natural Sciences

Citation: Ultrastructure of a condensed chromosome-like structure in a cyanobacterium (2016, October 21) retrieved 18 April 2024 from <https://phys.org/news/2016-10-ultrastructure-condensed-chromosome-like-cyanobacterium.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.