

# From unknown to beardog: Findings rescue fossils from 'trashbin' genus

October 12 2016

---



Artist's reconstruction of an early (ca. 38 million year-old) beardog from Texas, based on fossils of *Angelarctocyon australis* and *Gustafsonia cognita*. Credit: Credit: Monica Jurik, The Field Museum.

Large, ferocious-looking animals called beardedogs—neither bears nor dogs - roamed the northern hemisphere between about 40 and 5 million years ago.

But because so little data on their earliest members are available, their evolutionary relationships or phylogeny - and their place on the tree of life - has remained unclear.

A new study published today in *Royal Society Open Science* and based on improved phylogenetic analysis and advanced computed tomography (CT) scanning has changed that. The research identifies two fossils previously thought to be generic carnivorans (a large, diverse order of mammals) as some of the earliest known members of the beardedog family. These fossils are from animals estimated to be no larger than about five pounds, roughly the size of a Chihuahua and much smaller than formidable descendants that would later evolve.

The work reveals that while evidence of beardedogs has been found throughout the Northern Hemisphere, they may have originated or initially diversified in parts of what is now the southwestern U.S.

"Our research pinpoints the southwestern US as a key region in understanding the diversification and proliferation of this once successful group of predators prior to their extinction millions of years ago," said study coauthor Jack Tseng, PhD, assistant professor in the Department of Pathology and Anatomical Sciences in the Jacobs School of Medicine and Biomedical Sciences at the University at Buffalo.

## **The evolutionary roots of beardedogs**

First described back in 1986, fossils found in Texas of animals believed to be less than 5 pounds were originally assigned to the genus *Miacis*, a kind of "miscellaneous" category for early carnivores, based primarily

on external features.

"It was the best that could be done at the time," said Tseng, who did the work as a postdoctoral fellow at the American Museum of Natural History in New York. Susumu Tomiya, PhD, postdoctoral scholar at The Field Museum in Chicago, is lead author on the paper.

According to Tseng, the early anatomists based their hypothesis on superficial features like the shapes of the teeth and the locations of cranial openings where the blood vessels and nerves radiate from the brain and other external characteristics, assigning it to the genus *Miacis*.

"It's a kind of 'trashbin' genus, when the question is, well, what else could it be?" Tseng explained. "Now we've taken these fossils out of the trashbin and put them at the base of the beardog tree.

"We're not saying we've solved where they fit on the tree of life, but it's the most progress that's been made in quite awhile. Our work provides a clearer connection between the rest of the beardog family and their evolutionary roots."

## **A mysterious museum specimen**

Tomiya decided to study the fossils when he 'stumbled' upon one of the specimens in the Field Museum collection.

"I thought it looked odd and too advanced for what it had been claimed to be—a more primitive carnivore," said Tomiya. "It reminded me of some much larger beardogs so I decided to take a closer look."

That closer look included tapping Tseng's expertise with high resolution X ray CT 3D reconstructions of the intracranial anatomy of the fossils. While CT scans of the skull already existed, Tseng conducted a much

more detailed and time-consuming analysis through more than 1000 slices of CT scans of the skull. Adding to the difficulty of examining a very small skull—the whole animal was no larger than a Chihuahua—was the fact that the spaces inside the skull were still filled with rock.

Twenty years earlier, Tseng's PhD advisor, Xiaoming Wang of the Natural History Museum of Los Angeles County, had speculated that based on what could be seen externally, this animal was likely related to beardedogs, and may have had a deep embayment—a bone-enclosed space—in the region of its ear.

"The development of that feature is characteristic of beardedogs," said Tseng, "and it turns out that that actually is the case for the skull previously assigned to *Miacis*."

According to Tseng, the ear is very important in understanding mammalian evolution. "The ear can be used to calibrate how species are related," he said, noting that in other collaborative research, he is studying the ear's bony labyrinth, whose shape may be correlated to the kinds of movements an animal can make.

Tomiya's work conducting rigorous cladistic analyses (classification according to shared features) involved updating the taxonomy of these animals and their [evolutionary relationships](#) based on new phylogenetic affinities—in other words, how the beardedogs may be related to other carnivores that exist today, such as dogs, bears, raccoons and others.

**More information:** Whence the beardedogs? Reappraisal of the middle to late Eocene "Miacis" from Texas, USA, and the origin of Amphicyonidae (Mammalia, Carnivora), *Royal Society Open Science*, [rsos.royalsocietypublishing.org ... /10.1098/rsos.160518](https://rsos.royalsocietypublishing.org/.../10.1098/rsos.160518)

Provided by University at Buffalo

Citation: From unknown to beardog: Findings rescue fossils from 'trashbin' genus (2016, October 12) retrieved 25 April 2024 from <https://phys.org/news/2016-10-unknown-beardog-fossils-trashbin-genus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.