


# A record of ancient tectonic stress on Mars

November 3 2016

---



This part of Acheron Fossae was imaged by the High Resolution Stereo Camera on ESA's Mars Express on 4 May 2016 during orbit 15641. The image is centred on 36°N/142°W and the ground resolution is about 18 m per pixel. Credit: ESA/DLR/FU Berlin, CC BY-SA 3.0 IGO

Sets of ridges and troughs some 1000 km north of the giant Olympus Mons volcano contain a record of the intense tectonic stresses and strains experienced in the Acheron Fossae region on Mars 3.7–3.9 billion years ago.


This scene, captured by ESA's Mars Express on 4 May, focuses on the western part of Acheron Fossae, an isolated block of ancient terrain that covers an area about 800 km long and 280 km wide and stands up to 2 km higher than the surrounding plains.

Acheron Fossae is part of a network of fractures that radiates from the Tharsis 'bulge' some 1000 km to the south, home to the largest volcanoes on Mars. As the Tharsis region swelled with hot material rising from deep inside Mars as the volcanoes formed, it stretched and pulled apart the crust along lines of weakness over a wide area.

This process gave rise to the classic 'horst and graben' system – a series of depressions (graben) bounded by faults and uplifted blocks (horsts) either side of the graben.

The pattern of cross-cutting faults seen in various places in Acheron Fossae implies the region experienced stresses from different directions over time, suggesting a complex history.


Part of a dominant, curved ridge that extends through the entire region is seen in the lower left of the scene. It may be an ancient graben that has since been filled with material that has flowed along it, possibly from rock-laden glaciers that were deposited in more recent cold climatic conditions, long after the graben itself formed.



This context image shows part of Acheron Fossae on Mars imaged by Mars Express on 4 May 2016 during orbit 15641 (outlined by the large white box).  
Credit: NASA MGS MOLA Science Team


Acheron Fossae has been likened to Earth's continental rift systems. Major rift zones on Earth are associated with plate tectonics, such as mid-ocean ridges that are spreading apart.

On Mars, rifts are important for studies of the general evolution of the crust as well as the thermal evolution of the deeper subsurface.



The colour-coded topographic view shows relative heights and depths of terrain in the western portion of the Acheron Fossae region on Mars. Browns and whites represent the highest terrain (lower left), which transition through red, to yellow, green and blue in the upper right, representing progressively lower terrain. The image is based on a digital terrain model of the region, from which the topography of the landscape can be derived. The region was imaged by the High Resolution Stereo Camera on ESA's Mars Express on 4 May 2016 during orbit 15641. The image is centred on 36°N/142°W and the ground resolution is about 18 m per pixel. Credit: ESA/DLR/FU Berlin, CC BY-SA 3.0 IGO





This anaglyph image showing part of the Acheron Fossae region on Mars provides a 3D view of the landscape when viewed using stereoscopic glasses with red–green or red–blue filters. The image is based on data acquired by the nadir channel and one stereo channel of the High Resolution Stereo Camera on Mars Express on 4 May 2016 during orbit 15641. The main image is centred on 36°N/142°W and the ground resolution is about 18 m per pixel. Credit: ESA/DLR/FU Berlin, CC BY-SA 3.0 IGO

Provided by European Space Agency

Citation: A record of ancient tectonic stress on Mars (2016, November 3) retrieved 2 May 2024 from <https://phys.org/news/2016-11-ancient-tectonic-stress-mars.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.