

Researchers find biggest exposed fault on Earth

November 28 2016



Credit: Australian National University

Geologists have for the first time seen and documented the Banda Detachment fault in eastern Indonesia and worked out how it formed.

Lead researcher Dr Jonathan Pownall from The Australian National University (ANU) said the find will help researchers assess dangers of future tsunamis in the area, which is part of the Ring of Fire - an area



around the Pacific Ocean basin known for earthquakes and volcanic eruptions.

"The abyss has been known for 90 years but until now no one has been able to explain how it got so deep," Dr Pownall said.

"Our research found that a 7 km-deep abyss beneath the Banda Sea off eastern Indonesia was formed by extension along what might be Earth's largest-identified exposed <u>fault plane</u>."

By analysing high-resolution maps of the Banda Sea floor, geologists from ANU and Royal Holloway University of London found the rocks flooring the seas are cut by hundreds of straight parallel scars.

These wounds show that a piece of crust bigger than Belgium or Tasmania must have been ripped apart by 120 km of extension along a low-angle crack, or detachment <u>fault</u>, to form the present-day oceanfloor depression.

Dr Pownall said this fault, the Banda Detachment, represents a rip in the ocean floor exposed over 60,000 square kilometres.

"The discovery will help explain how one of the Earth's deepest sea areas became so deep," he said.





Credit: Australian National University

Professor Gordon Lister also from the ANU Research School of Earth Sciences said this was the first time the fault has been seen and documented by researchers.

"We had made a good argument for the existence of this fault we named the Banda Detachment based on the bathymetry data and on knowledge of the regional geology," said Professor Lister.

Dr Pownall said he was on a boat journey in eastern Indonesia in July when he noticed the prominent landforms consistent with surface extensions of the <u>fault line</u>.

"I was stunned to see the hypothesised fault plane, this time not on a computer screen, but poking above the waves," said Dr Pownall.



He said rocks immediately below the fault include those brought up from the mantle.

"This demonstrates the extreme amount of extension that must have taken place as the oceanic crust was thinned, in some places to zero," he said.

Dr Pownall also said the discovery of the Banda Detachment fault would help assesses dangers of future tsunamis and earthquakes.

"In a region of extreme tsunami risk, knowledge of major faults such as the Banda Detachment, which could make big earthquakes when they slip, is fundamental to being able to properly assess tectonic hazards," he said.

More information: Jonathan M. Pownall et al. Rolling open Earth's deepest forearc basin, *Geology* (2016). <u>DOI: 10.1130/G38051.1</u>

Provided by Australian National University

Citation: Researchers find biggest exposed fault on Earth (2016, November 28) retrieved 5 May 2024 from <u>https://phys.org/news/2016-11-biggest-exposed-fault-earth.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.