

Groundwater helium level could signal potential risk of earthquake

November 29 2016



Groundwater samples were obtained from seven locations in the Futagawa-Hinagu fault zones in the Kumamoto region. The locations are marked as yellow squares (denoted HRY, TMN, UKI, KKC, OTS, AJS, and MFN) on the map. The epicenters of the two tremors that preceded the main quake and the main



shock are shown in red circles. Credit: 2016 Yuji Sano.

Japanese researchers have revealed a relationship between helium levels in groundwater and the amount of stress exerted on inner rock layers of the earth, found at locations near the epicenter of the 2016 Kumamoto earthquake. Scientists hope the finding will lead to the development of a monitoring system that catches stress changes that could foreshadow a big earthquake.

Several studies, including some on the <u>massive earthquake</u> in Kobe, Japan, in 1995, have indicated that changes to the chemical makeup of groundwater may occur prior to earthquakes. However, researchers still needed to accumulate evidence to link the occurrence of earthquakes to such chemical changes before establishing a strong correlation between the two.

A team of researchers at the University of Tokyo and their collaborators found that when stress exerted on the earth's crust was high, the levels of a helium isotope, helium-4, released in the groundwater was also high at sites near the epicenter of the 2016 Kumamoto <u>earthquake</u>, a magnitude 7.3 quake in southwestern Japan, which caused 50 fatalities and serious damage.

The team used a submersible pump in deep wells to obtain groundwater samples at depths of 280 to 1,300 meters from seven locations in the fault zones surrounding the epicenter 11 days after the earthquake in April 2016. They compared the changes of helium-4 levels from chemical analyses of these samples with those from identical analyses performed in 2010.





Locations with high levels of stress exertion exhibited high levels of helium release, while locations with low levels of stress exertion exhibited low levels of helium release, thereby demonstrating a positive correlation. Credit: 2016 Yuji Sano.

"After careful analysis and calculations, we concluded that the levels of helium-4 had increased in samples that were collected near the epicenter due to the gas released by the rock fractures," says lead author Yuji



Sano, a professor at the University of Tokyo's Atmosphere Ocean Research Institute.

Furthermore, scientists estimated the amount of helium released by the rocks through rock fracture experiments in the laboratory using rock samples that were collected from around the earthquake region. They also calculated the amount of strain exerted at the sites for groundwater sample collection using satellite data. Combined, the researchers found a positive correlation between helium amounts in groundwater and the stress exertion, in which helium content was higher in areas near the epicenter, while concentrations fell further away from the most intense seismic activity.

"More studies should be conducted to verify our correlation in other earthquake areas," says Sano. "It is important to make on-site observations in studying earthquakes and other natural phenomena, as this approach provided us with invaluable insight in investigating the Kumamoto earthquake," he adds.





Scientists collect groundwater sample from a well in the Kumamoto earthquake region. Credit: 2016 Yuji Sano

More information: Yuji Sano, Naoto Takahata, Takanori Kagoshima, Tomo Shibata, Tetsuji Onoue & Dapeng Zhao, "Groundwater helium anomaly reflects strain change during the 2016 Kumamoto earthquake in Southwest Japan", *Scientific Reports* <u>dx.doi.org/10.1038/srep37939</u>

Provided by University of Tokyo



Citation: Groundwater helium level could signal potential risk of earthquake (2016, November 29) retrieved 2 May 2024 from <u>https://phys.org/news/2016-11-groundwater-helium-potential-earthquake.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.