

# Hot on the heels of quasiparticles

#### November 2 2016, by Oliver Morsch



A polaron (orange) is formed amidst the electrons (violet) inside a solid. Credit: ETH Zurich / Meinrad Sidler

Electrons in a solid can team up to form so-called quasiparticles, which lead to new phenomena. Physicists at ETH in Zurich have now studied previously unidentified quasiparticles in a new class of atomically thin semiconductors. The researchers use their results to correct a prevailing misinterpretation.



If one tries to understand weather phenomena, it's not much use looking at the behaviour of single water droplets or air molecules. Instead, meteorologists (and also laymen) speak of clouds, winds and precipitation - objects that result from the complex interplay between small particles. Physicists dealing with the optical properties or the conductivity of solids use much the same approach. Again, tiny particles - <u>electrons</u> and atoms - are responsible for a multitude of phenomena, but an illuminating picture only emerges when many of them are grouped into "quasiparticles".

However, finding out precisely what quasiparticles arise inside a material and how they influence one another is not a simple task, but more akin to a large puzzle whose pieces fit together, little by little, through arduous research. In a combination of experimental and <u>theoretical</u> <u>studies</u>, Ataç Imamoglu and his collaborators at the Institute for Quantum Electronics at the ETH in Zurich have now succeeded in finding a new piece of the puzzle, which also helps to put a previously misplaced piece in its correct position.

### **Excitons and polarons**

In solids quasiparticles can be created, for instance, when a photon is absorbed. The motional energy of electrons teeming about in a solid can only take values within well-defined ranges known as bands. A photon can promote an electron from a low-lying to a high-lying energy band, thus leaving behind a "hole" in the lower band.

The excited electron and the resulting hole attract each other through the electrostatic Coulomb force, and if that attraction is strong enough, the electron-hole pair can be viewed as a quasiparticle - an "exciton" is born. Two electrons and a hole can bind together to form a trion. When excitons and a large number of <u>free electrons</u> are simultaneously present however, the description of the qualitatively new - or "emergent" -



properties of the material requires the introduction of new type of quasiparticles called Fermi polarons.

# **Quasiparticles in a semiconductor**

Imamoglu and his colleagues wanted to find out the nature of quasiparticles that appear in a certain type of semiconductors in which electrons can only move in two dimensions. To do so, they took a single layer of molybdenum diselenide that is thousand times thinner than a micrometer and sandwiched it between two disks of boron nitride. They then added a layer of graphene in order to apply an electric voltage with which the density of electrons in the material could be controlled. Finally, everything was placed between two mirrors that formed an optical cavity.

With this complex experimental setup the physicists in Zurich could now study in detail how strongly the material absorbs light under different conditions. They found that when the semiconductor structure is optically excited, Fermi-polarons are formed - and not, as previously thought, excitons or trions. "So far, researchers - myself included - have misinterpreted the data available at the time in that respect", admits Imamoglu. "With our new experiments we are now able to rectify that picture."

#### Team effort with a guest scientist

"This was a team effort with essential contributions by Harvard professor Eugene Demler, who collaborated with us over several months when he was an ITS fellow", says Meinrad Sidler who is a doctoral student in Imamoglus group. Since 2013 the Institute for Theoretical Studies (ITS) of the ETH has endeavoured to foster interdisciplinary research at the intersection between mathematics, theoretical physics and



computer science. In particular, it wants to facilitate curiosity-driven research with the aim of finding the best ideas in unexpected places.

The study by Imamoglu and his colleagues, now published in *Nature Physics*, is a good example for how this principle can be successful. In his own research, Eugene Demler deals with ultracold atoms, studying how mixtures of bosonic and fermionic atoms behave. "His insight into polarons in atomic gases and solids have given our research important and interesting impulses, which we may not have come up with on our own", says Imamoglu.

## Light induced superconductivity

The insights they have gathered will most likely keep Imamoglu and his collaborators busy for some time to come, as the interactions between bosonic (such as excitons) and fermionic (electrons) particles are the topic of a large research project for which Imamoglu won an Advanced Grant of the European Research Council (ERC) last year, and is also supported by the National Centre of Competence in Research Quantum Science and Technology (NCCR QSIT). A better understanding of such mixtures would have important implications for basic research, but exciting applications also beckon. For instance, a key goal of the ERC project is the demonstration of control of superconductivity using lasers.

**More information:** Meinrad Sidler et al, Fermi polaron-polaritons in charge-tunable atomically thin semiconductors, *Nature Physics* (2016). DOI: 10.1038/nphys3949

Provided by ETH Zurich



Citation: Hot on the heels of quasiparticles (2016, November 2) retrieved 24 May 2024 from <u>https://phys.org/news/2016-11-hot-heels-quasiparticles.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.