


For platinum catalysts, tiny squeeze gives big boost in performance

November 24 2016



Bottom: Platinum atoms attached to layers of lithium cobalt oxide contract when electricity is applied, boosting platinum catalytic efficiency by 90 percent. Top: Removing electrons separates the atoms and lowers efficiency by 40 percent. (1 Å = 0.1 nanometer) Credit: Haotian Wang

A nanosize squeeze can significantly boost the performance of platinum

catalysts that help generate energy in fuel cells, according to a new study by Stanford scientists.

The team bonded a [platinum catalyst](#) to a thin material that expands and contracts as electrons move in and out, and found that squeezing the platinum a fraction of a nanometer nearly doubled its catalytic activity. The findings are published in the Nov. 25 issue of the journal *Science*.

"In this study, we present a new way to fine-tune [metal catalysts](#) at the atomic scale," said lead author [Haotian Wang](#), a former graduate student at Stanford now at Harvard University. "We found that ordinary battery materials can be used to control the activity of platinum and possibly for many other metal catalysts."

The new technique can be applied to a wide range of clean technologies, Wang said, including fuel cells that use platinum catalysts to generate energy, and platinum electrolyzers that split water into oxygen and [hydrogen fuel](#).

"Our tuning technique could make fuel cells more energy efficient and increase their power output," said co-author Yi Cui, a professor of materials science and engineering at Stanford and of photon science at the SLAC National Accelerator Laboratory. "It could also improve the hydrogen-generation efficiency of water splitters and enhance the production of other fuels and chemicals."


Electronic structure

Catalysts are used to make chemical reactions go faster while consuming less energy. The performance of a metal catalyst depends on its electronic structure - that is, how the electrons orbiting individual atoms are arranged.

"The electronic structure of a catalyst needs to match the molecule of interest in order to achieve the chemical reaction you want," Wang explained. "You can adjust the [electronic structure](#) of a catalyst by compressing the atoms or pulling them apart."

The Stanford team introduced a novel way to compress or separate the atoms by 5 percent, a mere 0.01 nanometer.

"That might not seem like much, but it's really a lot," Cui said. "How did we achieve that? It's really a marriage of battery research and catalysis."



Haotian Wang and his Stanford colleagues developed a new way to fine-tune platinum catalysts in fuel cells and other clean-energy technologies. Credit: L.A. Cicero/Stanford News Service

Experimental electrode

The study focused on lithium cobalt oxide, a material widely used in batteries for cellphones and other electronic devices. The researchers stacked several layers of lithium cobalt oxide together to form a battery-like electrode.

"Applying electricity removes lithium ions from the electrode, causing it to expand by 0.01 nanometer," Cui said. "When lithium is reinserted during the discharge phase, the electrode contracts to its original size."

For the experiment, the Stanford team added several layers of platinum to the lithium [cobalt oxide](#) electrode.

"Because platinum is bonded to the edge, it expands with the rest of the electrode when electricity is added and contracts during discharge," Cui said.

Performance

Separating the platinum layers a distance of 0.01 nanometer, or 5 percent, had a significant impact on performance, Wang said.

"We found that compression makes platinum much more active," he said. "We observed a 90 percent enhancement in the ability of [platinum](#) to reduce oxygen in water. This could improve the efficiency of hydrogen fuel cells."

Stretching the electrode by 5 percent had the opposite effect, suppressing oxygen production by 40 percent, Wang said.

"This is a dream experiment for a theorist," said study co-author Jens Nørskov, a professor of chemical engineering at Stanford's SUNCAT

Center for Interface Science and Catalysis. "We predicted theoretically some years ago that straining a catalyst can be used to control its performance, and here is the experiment to show that our theory works well."

"Our technology offers a very powerful way to controllably tune catalytic behavior," Cui added. "Now, mediocre catalysts can become good, and good catalysts can become excellent."

More information: "Direct and continuous strain control of catalysts with tunable battery electrode materials" *Science*, [science.sciencemag.org/cgi/doi ... 1126/science.aaf7680](https://science.sciencemag.org/cgi/doi/10.1126/science.aaf7680)

Provided by Stanford University

Citation: For platinum catalysts, tiny squeeze gives big boost in performance (2016, November 24) retrieved 18 April 2024 from <https://phys.org/news/2016-11-platinum-catalysts-tiny-big-boost.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.