


Scientists discover a key signal in intercellular communication

November 29 2016



The regulatory function of ISGylation in the secretion of nanovesicles to the external medium. Credit: CNIC

A team of scientists at the Centro Nacional de Investigaciones Cardiovasculares Carlos III (CNIC), led by Prof. Francisco Sánchez-Madrid, has characterized a cell signal that impedes intercellular

communication and could play a central role in biomedical strategies such as gene therapy, vaccine design, and immunotherapy. The study, published today in *Nature Communications*, characterizes a signal that impedes the secretion of nanovesicles called exosomes. Cells secrete exosomes as a means of intercellular communication; however, certain viruses can use exosomes as "Trojan horses" to facilitate their propagation and entry into neighboring cells.

The signal, called ISGylation, has in the past been viewed mainly as an antiviral signal, although some studies show that it can also be activated by other stimuli such as a lack of oxygen, aging or cancer. "In these situations, the secretion of exosomes, and therefore communication between cells, can be affected by this modification," explains Dr. Carolina Villarroya.

The research team is dedicated to decoding the processes that control exosome secretion and exploring potential biomedical applications. As Dr. Sánchez-Madrid explains, "as well as acting as messengers in intercellular communication, exosomes are potential tools for [gene therapy](#), [vaccine design](#) and immunotherapy." Several clinical trials are underway to assess new treatments using this approach.

The *Nature Communications* article describes how an antiviral signal activates the programmed degradation of proteins involved in exosome degradation. According to Villarroya, this signal "marks specific proteins located in endosomes, the place where exosomes are formed. This mark redirects these proteins toward the degradation pathway and impedes exosome secretion." Sánchez-Madrid points out that this newly identified mechanism, through which cells defend themselves against infection by activating the degradation of their own proteins, "could also be exploited by external agents for their propagation."

More information: Carolina Villarroya-Beltri et al, ISGylation

controls exosome secretion by promoting lysosomal degradation of MVB proteins, *Nature Communications* (2016). [DOI: 10.1038/ncomms13588](https://doi.org/10.1038/ncomms13588)

Provided by Centro Nacional de Investigaciones Cardiovasculares

Citation: Scientists discover a key signal in intercellular communication (2016, November 29)
retrieved 23 April 2024 from <https://phys.org/news/2016-11-scientists-key-intercellular.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--