

Detective work across dingo fence reveals new factor in woody shrub invasion

December 5 2016



A dusky hopping mouse which eats the seeds of woody shrubs in semi-arid Australia. Credit: Ben Moore

Extermination of dingoes and the consequent loss of small mammals - not just overgrazing by livestock - have led to a rapid spread of woody

shrubs across semi-arid Australia, a new study shows.

Re-introduction of small mammals to desert areas could help prevent further invasion by the woody weeds, the researchers suggest.

The study in outback NSW compared conditions on either side of the dingo fence - a two-metre tall, 5600-kilometre long fence erected more than a century ago to keep dingoes out of eastern Australia.

Dingoes are common on the western side of the fence, but rare on the other side, due to intensive control measures including poisoning, trapping and shooting over many decades.

"Woody shrubs pose a major problem for farmers and their abundance in semi-arid areas has risen dramatically in recent decades," says study senior author UNSW Associate Professor Mike Letnic.

"These weeds reduce the availability of feed for livestock and hinder access for mustering. But removing them is contentious, because they are classified as native vegetation."

To measure the encroachment of [woody shrubs](#) into grasslands, the researchers, led by PhD candidate Christopher Gordon from Western Sydney University, examined [aerial photographs](#) of far western NSW and eastern South Australia spanning a 50 year period.



A dingo in semi-arid Australia. Extermination of dingoes and consequent loss of small mammals have led to a rapid spread of woody shrubs across semi-arid Australia. Credit: Ben Moore

They assessed four sites in the Strzelecki Desert, two on either side of the dingo fence. The results are published in the *Journal of Animal Ecology*.

"The fence provides a rare, large-scale natural experiment to examine the effects that a top predator such as the dingo has on ecosystems," says Dr Gordon.

"We found that shrub cover had increased on the dingo-free side in both areas, one that had been used for livestock grazing and one that had not

been grazed since the 1970s. This suggests overgrazing by livestock may not be the main cause of shrub spread, as had been thought.

"There was also a negligible increase in shrub cover in comparable areas on the other side of the fence, which suggests dingoes help reduce shrub spread," he says.

This led the team to investigate the activity of small mammals and they found they were more common in areas where the shrubs were sparse. They also looked in detail at the feeding habits of one small mammal, the dusky hopping mouse, and found it consumed significant amounts of woody shrub seeds.

The rate of survival of shrub seedlings was greater in areas where dingoes were rare and the rate of removal of seeds was lower there.



The dingo fence -- a 2-meter tall, 5,600-kilometer long fence erected more than a century ago to keep dingoes out of eastern Australia. Credit: Ben Moore

"Taken together, our results suggest a cascading effect. A loss of dingoes allows foxes and feral cats to thrive and prey on the small mammals that eat the shrub seeds. Fewer [small mammals](#) means the shrubs can spread," says Gordon.

"This is the first study to link shrub encroachment to loss of an apex predator like the dingo, and it could explain shrub spread in many other areas around the globe.

"Rewilding initiatives that aim to conserve or re-establish small mammal populations could help stop the invasion of these pervasive weeds in Australia," he says.

The aerial photographs were collected by government agencies in 1948, 1972, 1983 and 1999. Shrub cover was 26-48 per cent greater in areas where [dingoes](#) were rare compared with areas where they were common.

Provided by University of New South Wales

Citation: Detective work across dingo fence reveals new factor in woody shrub invasion (2016, December 5) retrieved 1 May 2024 from <https://phys.org/news/2016-12-dingo-reveals-factor-woody-shrub.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--