


Gene editing yields tomatoes that flower and ripen weeks earlier

December 5 2016



Improving on a much-loved variety of cherry tomato (left), the CSHL team used gene editing to get ripe fruit two weeks earlier (right). These images were taken at the same point after planting. Credit: Lippman lab, CSHL

Using a simple and powerful genetic method to tweak genes native to two popular varieties of tomato plants, a team at Cold Spring Harbor Laboratory (CSHL) has devised a rapid method to make them flower and produce ripe fruit more than 2 weeks faster than commercial

breeders are currently able to do.

This means more plantings per growing season and thus higher yield. In this case, it also means that the plant can be grown in latitudes more northerly than currently possible - an important attribute as the earth's climate warms.

"Our work is a compelling demonstration of the power of gene editing - CRISPR technology - to rapidly improve yield traits in crop breeding," says CSHL Associate Professor Zachary Lippman, who led the research. Applications can go far beyond the tomato family, he says, to include many major food crops like maize, soybean, and wheat that so much of the world depends upon.

Lippman clarifies that the technique his team publishes today in *Nature Genetics* is about more than simply increasing yield. "It's really about creating a genetic toolkit that enables growers and breeders in a single generation to tweak the timing of flower production and thus yield, to help adapt our best varieties to grow in parts of the world where they don't currently thrive."

At the heart of the method are insights obtained by Lippman and colleagues, including plant scientists at the Boyce Thompson Institute in Ithaca, NY and in France led by Dr. José Jiménez-Gómez, about the evolution of the flowering process in many crops and their wild relatives as it relates to the length of the light period in a day. Genetic research revealed why today's cultivated tomato plant is not very sensitive to this variable compared to wild relatives from South America. Somehow, it does not much matter to domesticated plants whether they have 12 hours of daylight or 16 hours; they flower at virtually the same point after planting.

A well-known hormonal system regulates flowering time - and hence the

time when the plant will generate its first ripe fruit. The hormone florigen and a counteracting "anti-florigen" hormone called SP (for SELF PRUNING) act together, in yin-yang fashion, to, respectively, promote or delay flowering. In one phase of the newly reported research, the investigators studied a wild tomato species native to the Galapagos Islands - near the equator, with days and nights close to 12 hours year-round. They wanted to learn why, when grown in northern latitudes with very long summer days, this plant flowered very late in the season and produced few fruits.




Using gene editing to improve a prized cherry tomato variety (left) yields a bushier plant (right) that can be planted more densely, boosting yield while speeding up ripening by about two weeks. These plants were photographed an equal number of days after planting. Credit: Lippman lab, CSHL

The wild equatorial tomato, they learned, was extremely sensitive to daylight length. The longer the day, the longer the time to flowering, whereas "when you have a shorter light period, as in the plant's native habitat, they flower faster," Lippman says. This suggested there was a genetic change in tomato plants that occurred at some point before or during the domestication of wild tomato plants. Lippman suspects these changes likely had already occurred when the Spanish conquistador Cortez brought tomatoes to Europe from Mexico in the early 16th century, beginning the era of the plant's widespread adoption in mid-northern latitudes.

Lippman and colleagues traced the loss of day-length sensitivity in domesticated tomatoes to mutations in a gene called SP5G (SELF PRUNING 5G). It's a member of the same family of florigen and anti-florigen genes that were already known to regulate flowering time in tomato.

Growing the wild tomato plant from the Galapagos in greenhouses and fields in New York, Lippman and colleagues observed a sharp spike in the expression and activity of the anti-florigen hormone encoded by the SP5G gene, causing flowering to occur much later. In domesticated tomato plants, in contrast, that surge of anti-florigen is much weaker.



Inhibiting expression of the SP5G gene with CRISPR causes rapid flowering, regardless of day length. The modified plant is on the right, shown the same number of days after planting as the "unedited" wild type plant. Credit: Lippman lab, CSHL

The team's principal innovation - generating varieties of cherry and roma tomatoes that flower much earlier than the domesticated varieties on which they are based - arises from the observation that while domesticated plants are notably insensitive to day length, "there was some residual expression of the anti-florigen SP5G gene," Lippman says.

This led the team to employ the gene-editing tool CRISPR to induce tiny mutations in the SP5G gene. The aim was to inactivate the gene entirely such that it did not generate any anti-florigen protein at all.

When this tweaked version of SP5G was introduced to popular roma and

cherry tomato varieties, the plants flowered earlier, and thus made fruits that ripened earlier. Tweaking another anti-florigen gene that makes [tomato plants](#) grow in a dense, compact, shrub-like manner made the early-flowering varieties even more compact and early-yielding - a trait the team calls "double-determinate."

"What we've demonstrated here is fast-forward breeding," Lippman says. "Now we have a simple strategy to completely eliminate daylight sensitivity in elite inbred and hybrid plants that are already being cultivated. This could enable growers to expand their geographical range of cultivation, simply by using CRISPR to rapidly 'adapt' tomato and other crops to more northern latitudes, where summers have very long days and very short growing seasons."

More information: Variation in the flowering gene SELF PRUNING 5G promotes day-neutrality and early yield in tomato, *Nature Genetics*, [nature.com/articles/doi:10.1038/ng.3733](https://doi.org/10.1038/ng.3733)

Provided by Cold Spring Harbor Laboratory

Citation: Gene editing yields tomatoes that flower and ripen weeks earlier (2016, December 5) retrieved 10 April 2024 from <https://phys.org/news/2016-12-gene-yields-tomatoes-ripen-weeks.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--