


New method detects residual contaminants in ultra-pure helium gas

December 15 2016



Scientists used this radio frequency discharge source to measure the extremely low abundance of ^3He in samples of purified helium, the gas used in precision neutron lifetime experiments. Credit: US Department of Energy

The gas that makes balloons float is also vital to scientific experiments. In these experiments, natural helium (He) is purified, but it contains a tiny bit of a slightly different form of helium, known as the isotope ^3He . A sample can contain just one ^3He in every million helium atoms. That's too much for many experiments. Many experiments require ultra-pure

helium, with a ^3He component at least another million times smaller, or one in a trillion of the He atoms. Although techniques are believed to produce ultra-pure helium, until recently no experimental methods have confirmed that the amount of ^3He present in a sample is indeed that small. Now, scientists at the ATLAS facility at Argonne National Laboratory have used accelerator mass spectrometry (AMS) to precisely measure the very small concentrations of ^3He present.

Scientists need ultra-pure [helium](#) for a wide range of experiments. For example, they use ultra-pure helium to study the longevity and other properties of a free neutron. Free neutrons can provide insights into the formation of the universe and physics beyond the Standard Model, if measured accurately. To determine the purity of the helium for this study, the team demonstrated an approach that reaches a level of precision several orders of magnitude beyond that of any other technique. The team also found that measuring the amount of troublesome ^3He in purified helium samples intended for neutron studies suggest the need for significant experimental corrections, due to neutron absorption by the residual ^3He present.

Answering difficult scientific questions about the nature of the universe requires isotopically purified helium (^4He). The isotope ^3He can contaminate the helium. Accurately measuring the amount of ^3He requires determining the $^3\text{He}/^4\text{He}$ ratio at values well below those that can be achieved with standard mass spectroscopy techniques.

Accelerator mass spectrometry provides the only way to directly measure the ^3He content in purified helium samples at the level of sensitivity required for the neutron lifetime experiment, which seeks to determine how long a free neutron survives. Scientists used the ATLAS facility to demonstrate measurements of $^3\text{He}/^4\text{He}$ ratios as small as 10^{-14} , or 1 in 100,000,000,000,000. In this work, scientists tuned the ATLAS accelerator, which serves as an ultra-precise mass filter, with specialized carbon [ions](#). They scaled the accelerator components to $^3\text{He}^+$. To reduce

atmospheric ^3He contamination, the team produced the $^3\text{He}^+$ ions in a new radio frequency helium discharge source that reduced naturally occurring background sources of ^3He . They monitored the final accelerator tune by regularly switching to H^{3+} ions from high-purity hydrogen. They eliminated H_3^+ ions and ions consisting of paired deuterium and hydrogen atoms by dissociation in a gold foil, after acceleration to 8 MeV. After stripping the second electron from the $^3\text{He}^+$ ion, they dispersed the ions in a magnetic spectrograph and counted the $^3\text{He}^{2+}$ ions. The team anticipates that these observations will also guide the design of future neutron experiments. Based on known improvements, an ultimate sensitivity to $^3\text{He}/^4\text{He}$ ratios as small as 10^{-15} appears to be feasible.

More information: H. P. Mumm et al. High-sensitivity measurement of isotopic ratios for ultracold neutron experiments, *Physical Review C* (2016). [DOI: 10.1103/PhysRevC.93.065502](https://doi.org/10.1103/PhysRevC.93.065502)

Provided by US Department of Energy

Citation: New method detects residual contaminants in ultra-pure helium gas (2016, December 15) retrieved 26 June 2024 from <https://phys.org/news/2016-12-method-residual-contaminants-ultra-pure-helium.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.