


Mice can smell oxygen

December 2 2016



The green objects represent the type B cells - thus the neurons that are activated by low environmental oxygen. The red objects represent “canonical” or “conventional” olfactory sensory neurons: these cells each express one of the 1,100 odorant receptor genes in the genome, and respond to conventional odorous ligands. Credit: Max Planck Research Unit for Neurogenetics, Frankfurt am Main

The genome of mice harbours more than 1000 odorant receptor genes, which enable them to smell myriad odours in their surroundings. Researchers at the Max Planck Research Unit for Neurogenetics in Frankfurt, the University of Saarland in Homburg, the University of Cambridge and the Karolinska Institute in Stockholm have discovered that mice can also sense the oxygen level of the inhaled air using neurons in their nose. For this newly discovered sensory property, mice rely on two genes termed *Gucy1b2* and *Trpc2*, but apparently not on odorant receptor genes.

The research team discovered that a specific type of chemosensory neuron in the mouse [olfactory mucosa](#) responds to oxygen decreases in the environment. Chemosensory cells typically detect an increase in the concentration of a substance. In mammals, a lack of oxygen was thought to be detected primarily by the carotid body, a sensory organ situated at the carotid arteries in the neck. Activation of the carotid body results in activation of the respiratory centre in the brain. As mice live in burrows, it appears that during evolution an additional mechanism has developed in order to protect the individuals and their offspring from a shortage of oxygen.

The function of these so-called type B cells was enigmatic until this paper. "We activated these cells with low-oxygen air, and discovered a possibly vital function of these cells", says Peter Mombaerts, director of the Max Planck Research Unit for Neurogenetics. The research team investigated how type B cells behave when exposed to various oxygen levels. Using a calcium-sensitive dye, they observed that type B cells in the olfactory mucosa are activated following a modest decrease of the oxygen level in the external environment.

The scientists found that the *Gucy1b2* and *Trpc2* genes are essential for signal transduction in type B cells upon exposure to low oxygen. They used genetically altered mice in which either gene is inactivated

genetically. These genes encode, respectively, an enzyme that produces the second messenger cGMP and an ion channel through which calcium enters the cells. (Calcium is also another important intracellular messenger.) Without functional *Gucy1b2* and *Trpc2* genes, calcium-dependent signalling pathways are not activated in type B cells, and mice cannot distinguish or respond properly to decreased oxygen levels. Thus far the molecular sensor that detects the low oxygen level remains unknown. The researchers are investigating further the signalling mechanisms that result in activation of these neurons.

Moreover, the scientists found that mice can learn very quickly where locations with low oxygen levels are, and then avoid these areas. By contrast, mice with inactivated *Gucy1b2* or *Trpc2* genes cannot distinguish between normal and modestly decreased oxygen levels in the external environment, and do not show avoidance behaviour of these areas with a low oxygen level. These genes thus enable mice early on to select locations with an optimal oxygen level.

The researchers speculate that type B cells have also a social influence in mice. For instance, [mice](#) build their nests preferably in locations with a higher oxygen level in order to protect their offspring. "The offspring needs sufficient oxygen, otherwise they would be underserved", says Frank Zufall, director at the Center for Integrative Physiology and Molecular Medicine in Homburg.

Mice may be more sensitive to a lack of oxygen than humans. The human *Gucy1b2* and *Trpc2* genes are pseudogenes, which means that these [genes](#) may not encode proteins. It remains to be determined if type B [cells](#) occur in humans at all and if so, whether they detect low [oxygen levels](#).

More information: Katherin Bleymehl et al. A Sensor for Low Environmental Oxygen in the Mouse Main Olfactory Epithelium,

Neuron (2016). [DOI: 10.1016/j.neuron.2016.11.001](https://doi.org/10.1016/j.neuron.2016.11.001)

Provided by Max Planck Society

Citation: Mice can smell oxygen (2016, December 2) retrieved 19 April 2024 from <https://phys.org/news/2016-12-mice-oxygen.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.