


NASA sees wind shear's effects on Tropical Cyclone Yvette

December 23 2016



On Dec. 23 NASA-NOAA's Suomi NPP satellite captured this visible image of Tropical Cyclone Yvette north of Western Australia as it was being affected by vertical wind shear. Credit: NASA Goddard MODIS Rapid Response Team

Tropical Storm Yvette was being battered by vertical wind shear when

NASA-NOAA's Suomi NPP satellite passed over the Southern Indian Ocean.

On Dec. 23 the Visible Infrared Imaging Radiometer Suite (VIIRS) instrument aboard NASA-NOAA's Suomi NPP [satellite](#) provided a visible-light image of Tropical Cyclone Yvette that showed the bulk of clouds and thunderstorms were being pushed north and west of the center from [vertical wind shear](#). The Joint Typhoon Warning Center noted that [satellite imagery](#) showed a fully-exposed, well-defined low-level circulation center displaced east of isolated, flaring convection and developing thunderstorms.

On Dec. 23 at 4 a.m. EST (0900 GMT) Yvette's maximum sustained winds had decreased to 40 mph (35 knots/64.2 kph). It was centered near 14.7 degrees south latitude and 116.2 degrees east longitude, about 365 nautical miles northwest of Port Hedland, Australia. Yvette was moving east-southeast at 10.3 mph (9 knots/16.6 kph).

For forecast updates from Australia's Bureau of Meteorology, visit: <http://www.bom.gov.au/cyclone/index.shtml>.

The storm will accelerate and make landfall on Christmas Day (GMT) south of Broome, Western Australia.

Provided by NASA's Goddard Space Flight Center

Citation: NASA sees wind shear's effects on Tropical Cyclone Yvette (2016, December 23) retrieved 18 April 2024 from <https://phys.org/news/2016-12-nasa-effects-tropical-cyclone-yvette.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.