


NASA releases new eye-popping view of carbon dioxide

December 14 2016, by Alan Buis



Using observations from NASA's Orbiting Carbon Observatory-2, scientists have developed a new model of carbon behavior in our atmosphere from Sept. 1, 2014, to Aug. 31, 2015. Such models can be used to better understand and predict where carbon dioxide concentrations could be especially high or low. Credit: NASA Goddard Space Flight Center/K. Mersmann, M. Radcliff, producers

A new NASA supercomputer project builds on the agency's satellite

measurements of carbon dioxide and combines them with a sophisticated Earth system model to provide one of the most realistic views yet of how this critical greenhouse gas moves through the atmosphere.

Scientists have tracked the rising concentration of heat-trapping carbon dioxide for decades using ground-based sensors in a few places. A high-resolution visualization of the new combined data product provides an entirely different perspective. The visualization was generated by the Global Modeling and Assimilation Office at NASA's Goddard Space Flight Center, Greenbelt, Maryland, using data from the agency's Orbiting Carbon Observatory-2 (OCO-2) satellite, built and operated by NASA's Jet Propulsion Laboratory, Pasadena, California.

The 3-D visualization reveals in startling detail the complex patterns in which carbon dioxide in the atmosphere increases, decreases and moves around the globe over the time period from September 2014 to September 2015.

Atmospheric carbon dioxide acts as Earth's thermostat. Rising concentrations of the [greenhouse gas](#), due primarily to the burning of fossil fuels for energy, have driven Earth's current long-term warming trend. The visualization highlights the advances scientists are making in understanding the processes that control how much emitted carbon dioxide stays in the atmosphere and how long it stays there—questions that ultimately will determine Earth's future climate.

Scientists know that nearly half of all human-caused emissions are absorbed by the land and ocean. The current understanding is that about 50 percent of emissions remain in the atmosphere, about 25 percent are absorbed by vegetation on the land, and about 25 percent are absorbed by the ocean. However, those seemingly simple numbers leave scientists with critical and complex questions: Which ecosystems, especially on land, are absorbing what amounts of carbon dioxide? Perhaps most

significantly, as emissions keep rising, will the land and the ocean continue this rate of absorption, or reach a point of saturation?

The new dataset is a step toward answering those questions, explained Lesley Ott, a carbon cycle scientist at NASA Goddard and a member of the OCO-2 science team. Scientists need to understand the processes driving the "carbon flux"—the exchange of carbon dioxide between the atmosphere, land and ocean, Ott said.

"We can't measure the flux directly at high resolution across the entire globe," she said. "We are trying to build the tools needed to provide an accurate picture of what's happening in the atmosphere and translate that to an accurate picture of what's going on with the flux. There's still a long way to go, but this is a really important and necessary step in that chain of discoveries about carbon dioxide."

OCO-2, launched in 2014, is NASA's first satellite designed specifically to measure [atmospheric carbon dioxide](#) at regional scales.

"Since September of 2014, OCO-2 has been returning almost 100,000 carbon dioxide estimates over the globe each day. Modeling tools like those being developed by our colleagues in the Global Modeling and Assimilation Office are critical for analyzing and interpreting this high-resolution dataset," said David Crisp, OCO-2 science team leader at JPL.

The Global Modeling and Assimilation Office (GMAO) has previously included carbon dioxide in its GEOS Earth system model, which is used for all manner of atmospheric studies. This new product builds on that work by using the technique of data assimilation to combine the OCO-2 observations with the model. "Data assimilation is the process of blending model simulations with real-world measurements, with the precision, resolution and coverage needed to reflect our best understanding of the exchange of carbon dioxide between the surface

and atmosphere," explained Brad Weir, a researcher based in the GMAO.

The visualization showcases information about global carbon dioxide that has not been seen before in such detail: The rise and fall of carbon dioxide in the Northern Hemisphere throughout a year; the influence of continents, mountain ranges and ocean currents on weather patterns and therefore carbon dioxide movement; and the regional influence of highly active photosynthesis in places like the U.S. Corn Belt.

While the finely detailed carbon dioxide fluctuations are eye-catching, they also remind GMAO chief Steven Pawson of the progress scientists are making with computer models of the Earth system. One future step will be to integrate a more complex biology module into the model to better target the questions of [carbon dioxide](#) absorption and release by forests and other land ecosystems.

The results highlighted here demonstrate the value of NASA's unique capabilities in observing and modeling Earth. They also emphasize the collaboration among NASA centers and the value of powerful supercomputing. The assimilation was created using a model called the Goddard Earth Observing System Model-Version 5 (GEOS-5), which was run by the Discover supercomputer cluster at Goddard's NASA Center for Climate Simulation.

"It's taken us many years to pull it all together," Pawson said. "The level of detail included in this dataset gives us a lot of optimism that our models and observations are beginning to give a coherent view of the carbon cycle."

More information: For more information about OCO-2: oco.jpl.nasa.gov/

Provided by NASA

Citation: NASA releases new eye-popping view of carbon dioxide (2016, December 14)
retrieved 23 April 2024 from
<https://phys.org/news/2016-12-nasa-eye-popping-view-carbon-dioxide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.