


Partnership at a distance: Deep-frozen helium molecules

December 7 2016



WERDEN ZWEI ÜBERZEUGTE EINZELGÄNGER
IN EIN GEMEINSAMES BETT GEZWUNGEN, RÜCKEN SIE
BIS ÜBER DEN RAND HINAUS VONEINANDER AB.


"When two loners are forced to share a bed, they move well beyond its edges to get away from each other." Credit: Peter Evers

Helium atoms are loners. Only if they are cooled down to an extremely low temperature do they form a very weakly bound molecule. In so doing, they can keep a tremendous distance from each other thanks to the quantum-mechanical tunnel effect. As atomic physicists in Frankfurt have now been able to confirm, over 75 percent of the time they are so far apart that their bond can be explained only by the quantum-mechanical tunnel effect.

The [binding energy](#) in the helium molecule amounts to only about a billionth of the binding energy in everyday molecules such as oxygen or nitrogen. In addition, the molecule is so huge that small viruses or soot particles could fly between the atoms. This is due, physicists explain, to the quantum-mechanical "tunnel effect". They use a potential well to illustrate the bond in a conventional molecule. The atoms cannot move further away from each other than the "walls" of this well. However, in quantum mechanics the atoms can tunnel into the walls. "It's as if two people each dig a tunnel on their own side with no exit", explains Professor Reinhard Dörner of the Institute of Nuclear Physics at Goethe University Frankfurt.

Dörner's research group has produced this helium molecule in the laboratory and studied it with the help of the COLTRIMS reaction microscope developed at the University. The researchers were able to determine the strength of the bond with a level of precision not previously achieved and measured the distance between the two atoms in the molecule. "The helium molecule is something of a touchstone for quantum-mechanical theories, as the value of the binding energy theoretically predicted is heavily dependent on how accurately all physical and quantum-mechanical effects were taken into account", explains Dörner.

Even the theory of relativity, which is otherwise mainly required for astronomical calculations, had to be incorporated here. "If even just a small mistake occurs, the calculations produce major deviations or even indicate that a helium molecule cannot exist at all", says Dörner. The precision measurements performed by his research group will serve as a benchmark for future experiments.



Experimental setup: The helium molecules are produced in a gas jet at an extremely low temperature and separated from the rest of the gas jet in a diffraction grating. The X-ray flash (red) of the FLASH laser (Hamburg) ionises both helium atoms of the molecule so that they fly apart with explosive force. The ions are then mapped on a spatial resolution detector, as symbolized by the film strip. Credit: AG Dörner

Two years spent taking measurements in the cellar

Dörner's research group began investigating the helium molecule back in 2009, when the German Research Foundation awarded him a Reinhart Koselleck Project and funding to the tune of € 1.25 million. "This type of funding is risk capital, as it were, with which the German Research Foundation supports experiments with a long lead time", explains Dörner. He was thus able to design and set up the first experiments with his group. Initial results were achieved by Dr. Jörg Voigtsberger in the framework of his doctoral dissertation. "In the search for atoms which 'live in the tunnel', Jörg Voigtsberger spent two years of his life in the cellar", recalls Dr. Till Jahnke, senior lecturer and Voigtberger's supervisor at the time. It is there, in the cellar, that the laser laboratory of the atomic physics group is housed.

Stefan Zeller, the next doctoral researcher, considerably improved the equipment with the help of Dr. Maksim Kunitski and increased measurement precision still further. To do so, one of his tasks was to shoot at the very weakly bonded helium molecule with FLASH, the free-electron laser at the DESY research centre in Hamburg and the largest "photon canon" in Germany. "Stefan Zeller's work was remarkable. It was his untiring effort, his excellent experimental research skills and his ability not to be disheartened by temporary setbacks which made our success possible at all", remarks Professor Dörner, Zeller's doctoral supervisor.

Already beforehand the results have attracted considerable interest at national and international level. They will now appear in the renowned journal *Proceedings of the National Academy of Sciences (PNAS)* and are also part of the research work for which the group was awarded the Helmholtz Prize 2016.

More information: Stefan Zeller et al, Imaging the Hequantum halo state using a free electron laser, *Proceedings of the National Academy of Sciences* (2016). [DOI: 10.1073/pnas.1610688113](https://doi.org/10.1073/pnas.1610688113)

Provided by Goethe University Frankfurt am Main

Citation: Partnership at a distance: Deep-frozen helium molecules (2016, December 7) retrieved 18 April 2024 from

<https://phys.org/news/2016-12-partnership-distance-deep-frozen-helium-molecules.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.