


Physicists measure the loss of dark matter since the birth of the universe

December 28 2016



The discrepancy between the cosmological parameters in the modern Universe and the Universe shortly after the Big Bang can be explained by the fact that the proportion of dark matter has decreased. The authors of the study could calculate how much dark matter could have been lost and what the corresponding size of the unstable component would be. Researchers may explore how quickly this unstable part decays and say if dark matter is still disintegrating. Credit: MIPT

Russian scientists have discovered that the proportion of unstable particles in the composition of dark matter in the days immediately

following the Big Bang was no more than 2 percent to 5 percent. Their study has been published in *Physical Review D*.


"The discrepancy between the cosmological parameters in the modern [universe](#) and the universe shortly after the Big Bang can be explained by the fact that the proportion of dark matter has decreased. We have now, for the first time, been able to calculate how much dark matter could have been lost, and what the corresponding size of the unstable component would be," says co-author Igor Tkachev of the Department of Experimental Physics at INR.

Astronomers first suspected that there was a large proportion of hidden mass in the universe back in the 1930s, when Fritz Zwicky discovered "peculiarities" in a cluster of galaxies in the constellation Coma Berenices—the galaxies moved as if they were under the effect of gravity from an unseen source. This hidden mass, which is only deduced from its gravitational effect, was given the name dark matter. According to data from the Planck space telescope, the proportion of dark matter in the universe is 26.8 percent; the rest is "ordinary" matter (4.9 percent) and dark energy (68.3 percent).

The nature of dark matter remains unknown. However, its properties could potentially help scientists to solve a problem that arose after studying observations from the Planck telescope. This device accurately measured the fluctuations in the temperature of the [cosmic microwave background](#) radiation—the "echo" of the Big Bang. By measuring these fluctuations, the researchers were able to calculate key cosmological parameters using observations of the universe in the recombination era—approximately 300,000 years after the Big Bang.

However, when researchers directly measured the speed of the expansion of galaxies in the modern universe, it turned out that some of these parameters varied significantly—namely the Hubble parameter,

which describes the rate of expansion of the universe, and also the parameter associated with the number of galaxies in clusters. "This variance was significantly more than margins of error and systematic errors known to us. Therefore, we are either dealing with some kind of unknown error, or the composition of the ancient universe is considerably different to the modern universe," says Tkachev.



The concentration of the unstable component of dark matter F against the speed of expansion of non-gravitationally bound objects (proportional to the age of the

Universe) when examining various combinations of Planck data for several different cosmological phenomena. Credit: MIPT

The discrepancy can be explained by the decaying dark matter (DDM) hypothesis, which states that in the early universe, there was more dark matter, but then part of it decayed.

"Let us imagine that dark matter consists of several components, as in [ordinary matter](#) (protons, electrons, neutrons, neutrinos, photons). And one component consists of [unstable particles](#) with a rather long lifespan. In the era of the formation of hydrogen, hundreds of thousands of years after the Big Bang, they are still in the universe, but by now (billions of years later), they have disappeared, having decayed into neutrinos or hypothetical relativistic particles. In that case, the amount of dark matter in the era of hydrogen formation and today will be different," says lead author Dmitry Gorbunov, a professor at MIPT and staff member at INR.

The authors of the study analyzed Planck data and compared them with the DDM model and the standard Λ CDM (Lambda-cold dark matter) model with stable dark matter. The comparison showed that the DDM model is more consistent with the observational data. However, the researchers found that the effect of gravitational lensing (the distortion of cosmic microwave background radiation by a gravitational field) greatly limits the proportion of decaying dark matter in the DDM model.

Using data from observations of various cosmological effects, the researchers were able to give an estimate of the relative concentration of the decaying components of dark matter in the region of 2 percent to 5 percent.

"This means that in today's universe, there is 5 percent less dark matter

than in the recombination era. We are not currently able to say how quickly this unstable part decayed; [dark matter](#) may still be disintegrating even now, although that would be a different and considerably more complex model," says Tkachev.

More information: A. Chudaykin et al, Dark matter component decaying after recombination: Lensing constraints with Planck data, *Physical Review D* (2016). [DOI: 10.1103/PhysRevD.94.023528](https://doi.org/10.1103/PhysRevD.94.023528)

Provided by Moscow Institute of Physics and Technology

Citation: Physicists measure the loss of dark matter since the birth of the universe (2016, December 28) retrieved 23 April 2024 from <https://phys.org/news/2016-12-physicists-loss-dark-birth-universe.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--