

Predator invasion had devastating, long-term effects on native fish

December 12 2016



Peacock bass, *Cichla monoculus*. Credit: Diana Sharpe

In 1969, 60 to 100 peacock bass imported from Buga, Colombia, were introduced into a pond in Panama for sport fishing. Several individuals escaped. By the early 1970s, they colonized Gatun Lake, the reservoir forming the main channel of the Panama Canal. Forty-five years later, native fish populations in the lake still have not recovered according to a new Smithsonian report.

Peacock bass, *Cichla monoculus*, originally from the Amazon River and its tributaries, are voracious predators. They are considered delicious game [fish](#) and have been intentionally stocked for recreational and commercial fishing around the world.

In 1973, Thomas Zaret and Robert Paine, working at the Smithsonian Tropical Research Institute (STRI) field station on Barro Colorado Island, published a study in *Science* magazine showing that 60 percent of the native freshwater fish in Gatun Lake were extirpated after the accidental release. Their study was one of the first to demonstrate how devastating predator introductions can be and remains as one of the most highly cited studies on this subject.

"Very few studies have examined the long-term outcome of an invasion by a predatory fish," said Diana Sharpe, postdoctoral fellow in staff scientist Mark Torchin's lab at STRI and at Canada's McGill University. "This is the perfect place to test Zaret and Paine's prediction that certain species would never return to the [lake](#) by repeating their study 45 years later."



Diana Sharpe and co-author Rigoberto Gonzalez traveling to a study site on Lake Alajuela in Panama. Credit: Diana Sharpe

"Bob Paine passed away earlier this year, in his 80s," Sharpe said. "I had the opportunity to correspond with him just a few months before his death, and he was very excited that someone was finally doing a follow-up. He kindly Fed-Exed a huge stack of unpublished data related to their original work...all the way to Panama!"

Sharpe teamed up with two Panamanian scientists, Luis Fernando De León, assistant professor at the University of Massachusetts and research associate at INDICASAT-AIP in Panama, and Rigoberto González of

STRI to capture fish in gill nets and beach seines. They photographed, weighed, measured and released most of the fish. Several fish of each species were preserved for reference.

"Seining was very tricky because it involves wading through the shallows dragging a heavy net," Sharpe said. "The crocodile population in Gatun Lake has increased sharply since the 1970s. We had to abort our seining on more than one occasion after spotting crocs basking in the shallows. At one point, we actually caught a small croc and had to cut the net to release it."

"Zaret and Paine recorded 12 [native fish species](#) in the Trinidad arm of Gatun Lake in 1972 before the peacock bass invasion reached that region of the lake," Sharpe said. "We recaptured only three of those species after extensive seining. Our study shows that despite the appeal of peacock bass as a sport fish, the introduction of a novel apex predator can have dramatic and long-lasting impacts on native communities, even in diverse tropical communities, which are sometimes thought to be more resistant to invasion."

fish communities in two invaded lakes, Gatun and Alajuela, with fish communities in Lake Bayano, where there are no peacock bass. Gatun and Alajuela were also invaded by the jaguar cichlid, sometime between 2004 and 2013. The abundance of [native fish](#) was much greater in Lake Bayano than in the lakes containing peacock bass, corroborating the pattern observed in Gatun Lake through time.

This is not the first case of an introduced fish doing away with the natives. Nile perch, *Lates niloticus*, released into Lake Victoria in Africa caused the presumed extinction of 250 endemic cichlid fish in the 1980s.

"This study illustrates the value of historical and long-term data collected at the Smithsonian in Panama," Torchin said. "The comparison would not have been possible without Zaret and Paine's work 45 years ago around Barro Colorado. We never know when we're going to go back and use data from studies in the past, but in Panama we are lucky to have data on fish populations that goes back to Hildebrand and Meek's studies as part of the Smithsonian's Panama Biological Survey in 1910."

More information: D.M.T Sharpe et al, Tropical fish community does not recover 45 years after predator introduction, *Ecology* (2016). [DOI: 10.1002/ecy.1648](https://doi.org/10.1002/ecy.1648)

Provided by Smithsonian Tropical Research Institute

Citation: Predator invasion had devastating, long-term effects on native fish (2016, December 12) retrieved 27 April 2024 from <https://phys.org/news/2016-12-predator-invasion-devastating-long-term-effects.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.