

# Scientists zero in on biological diversity in 'poor man's rainforest'

December 20 2016

---



Michael Strickland, an assistant professor of biological sciences at Virginia Tech, sorts through a soil sample obtained from Pandapas Pond. Strickland is affiliated with the university's Global Change Center. Credit: Lindsay Key

The soil beneath our feet is not as biologically diverse as scientists

previously thought, according to a research team that includes a Virginia Tech soil microbial ecologist.

Leftover DNA from dead organisms—known as "relic DNA"—has historically thrown a wrench into estimates, causing scientists to overestimate [microbial diversity](#) by as much as 55 percent.

Understanding microbial diversity in soil is crucial for understanding how environmental processes like atmospheric nitrogen fixation and climate change occur.

But a team that includes Michael Strickland, an assistant professor of biological sciences in the College of Science, used a [high throughput sequencing](#) technique to determine the exact make-up of 31 soil samples from varying climates and ecosystems.

The results were published in *Nature Microbiology* this week.

"When we started to realize that our numbers could be off, we knew we had to find a way to take a closer look at how many species are actually there," said Strickland, who is also affiliated with the university's Global Change Center and the Fralin Life Science Institute.

Information about populations of [microbes](#) in soil is important because these organisms play critical roles in the terrestrial ecosystem and they help maintain [soil fertility](#).

But linking the activities of microbes to soil processes is difficult. Scientists need to measure living microbes—a challenging task because DNA from dead microbes can persist in soil for years, obscuring the analysis of microbial diversity.

"This research suggests that a significant proportion of the microorganisms detected in soil using DNA based techniques are no

longer living," said Ember Morrissey, an assistant professor of plant and soil sciences at West Virginia University who was not involved in the research project. "As a consequence we may need to use tools that distinguish the genetic material of living cells from the relic DNA of dead microbes in order to understand the influence of microbial 'species' on important ecosystem processes."

Paul Carini, a microbial ecophysiologicalist at the University of Colorado Boulder and first author of the paper, used PMA, a photoreactive dye that binds to relic DNA but does not adhere well to living cells, to distinguish viable cells from DNA debris in soil.

"Accounting for relic DNA in our analyses will help us understand the important ebb and flow of the soil microbiome and help us better understand how microbes help regulate soil fertility and make earth habitable in the face of a changing climate," said Carini.

Although [soil microbial communities](#) were found to be less diverse than previously thought, they are still pretty diverse, according to Strickland. In one gram of soil, thousands of species of microbes live, causing Strickland to deem [soil](#) as "the poor man's rainforest."

**More information:** Paul Carini et al. Relic DNA is abundant in soil and obscures estimates of soil microbial diversity, *Nature Microbiology* (2016). [DOI: 10.1038/nmicrobiol.2016.242](https://doi.org/10.1038/nmicrobiol.2016.242)

Provided by Virginia Tech

Citation: Scientists zero in on biological diversity in 'poor man's rainforest' (2016, December 20) retrieved 19 April 2024 from <https://phys.org/news/2016-12-scientists-biological-diversity-poor-rainforest.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.