

## Astronomers detect a strong outburst of an embedded young stellar object





The images of UKIDSS-J185318.36+012454.5 in different wavelengths: Ks band from Varricatt et al (2010) (left top), Ks band from DR6 UKIDSS GPS (right top), [3.6]  $\mu$ m band from GLIMPSE I survey (left bottom) and [5.8]  $\mu$ m band from GLIMPSE I survey (right bottom). The position of IRAS



18507+0121 source from IRAS PSC and IRAS PSC/FSC Combined Catalogue (Abrahamyan et al., 2015) marked by black and white crosses respectively. On DR6 UKIDSS GPS Ks image are marked three stellar sources (#43, 49 and 54, Shepherd et al., 2004). Credit: Nikoghosyan et al., 2016.

(Phys.org)—Astronomers from the Byurakan Astrophysical Observatory in Armenia have discovered a strong outburst of the embedded young stellar object (YSO) designated UKIDSS-J185318.36+012454.5. This newly detected event could provide important insights for the understanding of eruptive variability of young stars. The researchers presented their discovery in a paper published Dec. 30 on the arXiv preprint server.

UKIDSS-J185318.36+012454.5 is located in the vicinity of the IRAS 18507+0121 source and probably belongs to the massive star-forming region associated with GAL 034.4+00.23 HII region. The object is at least 5,000 light years away from the Earth and its neighborhood contains massive protostars with an age of about 100,000 years, as well as a low-mass stellar population approximately 1 million years old. The stellar object was classified as an intermediate mass YSO less than 100,000 years old.

Recently, a team of researchers led by Elena Nikoghosyan has combed through the data and images provided by the United Kingdom Infrared Telescope (UKIRT) in Hawaii to search for evidence of outbursts in starforming regions. They used the archival infrared and submillimeter data from UKIRT Galactic Plane Survey (UKIDSS GPS) and found that an <u>outburst</u> occurred in UKIDSS-J185318.36+012454.5 in the period of 2004-2006.

"UKIRT Galactic Plane Survey is a deep survey with good space



resolution. Therefore, it provides very useful information for researchers who study star forming regions. In particular, for detection of UKIDSS-J185318.36+012454.5 outburst, UKIRT Galactic Plane Survey was the basic source of information," Nikoghosyan told Phys.org.

According to the paper, the amplitude of brightness of the detected outburst is at least five mag. The object remained at this level of brightness until 2014.

The results of the research allowed the scientists to re-classify UKIDSS-J185318.36+012454.5 as an eruptive variable with mixed characteristics, namely a MNor-type object, and Nikoghosyan emphasized the significance of the study.

"Eruptions of pre-main sequence stellar objects are rare events, thus a new outburst is always noteworthy. And I think that the discovery of a new eruptive variable UKIDSS-J185318.36+012454.5 with a so strong outburst (delta K > 5 mag) was a big success," she said.

The authors of the paper also sought the most plausible explanation for eruptive variability of young stars. They noted that several physical mechanisms, such as rotation, cool or hot spots, accretion-driven wind and outflow can explain the near-infrared variability, but these mechanisms often produce short-term variability with amplitude that is not expected to exceed one mag in the K band.

"In our opinion, all of the proposed explanations of this phenomena to date (eruptive variability of <u>young stars</u>) do not explain it completely, most likely due to lack of information. With respect to this object, we can only assume that this outburst corresponds to the transition between two evolution stages: Class 0 (no optical and near-infrared emission) and Class I (generally optically obscured). But this is only an assumption," Nikoghosyan concluded.



The researchers now plan further observations of UKIDSS-J185318.36+012454.5. However, these studies could only be conducted in collaboration with scientists from other countries due to the fact that in Armenia, there is no opportunity to observe this object in the infrared range.

**More information:** New eruptive variable in the massive star-forming region associated with IRAS 18507+0121 source, arXiv:1612.09190 [astro-ph.SR] <u>arxiv.org/abs/1612.09190</u>

## Abstract

Aims. We report the discovery of a strong outburst of the embedded young stellar object (YSO), namely UKIDSS-J185318.36+012454.5, located in the star-forming region associated with IRAS 18507+0121 source and GAL 034.4+00.23 HII region. Methods. Using the archival photometric data and images we determined the amplitude and the epoch of the outburst, as well as the evolution stage and the basic parameters of the object. Results. According to the near and mid-infrared colors and spectral energy distribution, we classify the object as an intermediatemass young stellar object (YSO) with Class 0/I evolution stage. The outburst occurred in the period of 2004-2006. The amplitude of brightness is as least Ks = 5.0 mag. The summation of the photometric and spectral data does not allow to classify UKIDSS-J185318.36+012454.5 as FUor or EXor. We can consider it as an eruptive variable with mixed characteristics or MNor type object.

## © 2017 Phys.org

Citation: Astronomers detect a strong outburst of an embedded young stellar object (2017, January 3) retrieved 7 May 2024 from <u>https://phys.org/news/2017-01-astronomers-strong-outburst-embedded-young.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.