

# Integrated management of stink bugs in beans

January 4 2017

---



Credit: NIBIO – Norwegian Institute of Bioeconomy Research

In many tropical countries, stink bug infestations are a severe problem in common bean production. A new study looks at an integrated management system to control this agricultural pest.

Stink bugs are found throughout the world's temperate and tropical

areas. It is a serious agricultural pest and causes damage to a range of crops from common beans, soybeans, lima beans and corn to tobacco, peaches, grapes, tomatoes and cotton, as well as several species of weeds.

For common beans—an important protein crop and food staple in many countries in Latin America and sub-Saharan Africa—[stink bugs](#) present a major challenge.

In a new PhD thesis study, Yordanys Ramos González examines the biology, diversity, prevalence and management of stink bugs in common beans in Cuba to find innovative and practical solutions that may be used in integrated pest management (IPM) of these important pests in beans. The aim is to enable increased food production in Cuba and other tropical countries facing problems with stinkbugs in beans.

## **Important pests**

"The common bean is an important nutrient source in many countries, especially in Eastern and Southern Africa and Central and South America. In these parts of the world, the human diet lacks animal protein and is largely based on legumes to supply protein. The stink bug species are important pests of legumes in these areas," explains Ramos González.

According to the PhD student, the bug is also moving northwards, following the current climatic warming.

"Recently, breeding populations have been confirmed in Switzerland and Canada, but the pathways of entry and genetic diversity of most of these stink bugs remains unknown," he says.

Wild host plants play an important role as alternative host plants and may

provide good conditions for the build-up of stink bug populations.

Stink bugs are known to cause direct injury by puncturing the seeds and pods of legumes with their piercing-sucking mouthparts, preferentially young pods and developing seeds. Seed feeding can cause seed abortion or deformation. A secondary impact of stink bug damage is that the feeding provides an opportunity for plant pathogens such as bacteria and yeasts to colonize the plant.

## Using natural enemies



Nezara infested with beauveria. Credit: NIBIO

Several studies have been conducted on the biology, ecology and abundance of stink bugs in soybeans, but there is a lack of studies in common beans, and those that exist focus on the feeding behavior of stink bugs.

"Our aim was therefore to determine the seasonal population density fluctuation of stink bug species in the common bean, and evaluate the effect of three different bean cultivars on prevalence and damage caused by stink bugs.

"We also looked at organic versus conventional production and the natural occurrence of the beneficial fungus *Beauveria* that is a natural enemy of stink bugs. The thesis then evaluates the potential of using IPM strategies such as the seeding date and natural enemies such as insect pathogenic fungi and parasitoids in biological control of these stink bugs," Ramos González explains.

A field experiment was conducted in Cuba during two bean growing seasons (2010-2011 and 2012-2013) at three seeding dates to identify stink bug species and determine their seasonal population density fluctuation and damage caused in three cultivars.

In addition to conducting his field and laboratory work in Cuba, Ramos González have been on a yearly exchange to NIBIO to conduct molecular laboratory work, literature studies and write his PhD thesis papers.

## **Control with fungus**

Ramos González says the research results show that the main stink bugs found in bean agroecosystems in Cuba are the types *Nezara viridula* and *Piezdorus guildinii*. The researchers discovered that the populations peak at the time of fruit development.

"Our research shows that the bean cultivar ICA Pijao is the best choice to avoid stink bug damage," he says.

All the isolates of the beneficial fungus *Beauveria* obtained from the bean agroecosystems were molecularly identified as the species *Beauveria bassiana*. There was a higher occurrence of this beneficial fungus in organic than conventional agroecosystems.

He explains that the research showed that this fungus can be used to efficiently control stink bugs:

"Our study also evaluated the efficacy of *B. bassiana* in the control of stink bugs and we have found an isolate that is a promising biocontrol candidate for the use in integrated management in Cuba and elsewhere in the tropics," Ramos González says.

Provided by NIBIO – Norwegian Institute of Bioeconomy Research

Citation: Integrated management of stink bugs in beans (2017, January 4) retrieved 14 May 2024 from <https://phys.org/news/2017-01-bugs-beans.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--