

Finding may offer farmers a way to reduce harmful emissions from fertilized soil

January 16 2017, by David Malmquist



B.K. Song collects water samples from an over-fertilized lake for analyses of microbial communities and nutrients. Credit: David Malmquist

Those concerned with the health of Chesapeake Bay are familiar with nitrogen as a major pollutant whose excess runoff into bay waters can lead to algal blooms and low-oxygen dead zones. Perhaps less familiar is the significant role that a form of nitrogen gas plays in greenhouse warming and the destruction of Earth's ozone layer.

Now, an international group of scientists including B.K. Song of William & Mary's Virginia Institute of Marine Science has discovered that



production of this potent greenhouse gas—known as N2O or <u>nitrous</u> <u>oxide</u>—can be bypassed as complex nitrogen compounds in soil, water and fertilizers break down into the unreactive nitrogen gas (N2) that makes up most of our atmosphere.

Their discovery, published in a recent edition of *Scientific Reports*, reveals an entirely new pathway in the <u>global nitrogen cycle</u> and could lead to new ways for farmers and others to reduce their emissions of harmful gases. The study's lead author is Rebecca Phillips of New Zealand's Landcare Research Institute, along with Landcare colleagues Andrew McMillan, Gwen Grelet, Bevan Weir and Palmada Thilak. Also contributing to the study was Craig Tobias of the University of Connecticut.

Agriculture contributes more nitrous oxide to the atmosphere than any other human activity—primarily through <u>nitrogen fertilization</u>. This greenhouse gas is 300 times more effective at trapping heat than carbon dioxide and 10 times more effective than methane. Nitrous oxide also moves into the stratosphere and destroys ozone.





Associate Professor B.K. Song collects water samples from a local stream for analyses of nitrogen and microbial communities. Credit: David Malmquist

Current wisdom holds that nitrous oxide is inevitably produced when soil nitrogen—including fertilizer components such as ammonia, ammonium, and urea—breaks down. It's also thought this breakdown process, known as denitrification, requires the action of microbes and can only occur in the absence of oxygen.

The current research contradicts each of these long-held ideas.

"Our findings question the assumption that nitrous oxide is an intermediate required for formation of <u>nitrogen gas</u>," says Phillips. "They also throw doubt on whether microbial production of nitrous oxide must take place in the absence of oxygen."



A farmer applies nitrogen fertilizer to a field. Credit: The College of William & Mary

"We now have a pathway that doesn't require microbes," adds Song.



"The process of denitrification can happen abiotically, without the need for bacteria or fungi."

The team's discovery could lead to practical applications for decreasing the impacts of <u>excess nitrogen</u> in the environment, a topic they focused on while presenting their findings during a recent meeting in Washington, D.C., sponsored by the U.S. Department of Agriculture and the National Integrated Water Quality Program.

"It might give us a way to engineer the system to reduce levels of fixed nitrogen," says Song. "By changing the types and ratios of nitrogen compounds in fertilizer, you might have a better way to reduce excess nitrogen, and to mitigate eutrophication or nutrient enrichment in nearby waters."



Fertilization of home yards and gardens is another source of nitrogen to the



environment. Credit: The College of William & Mary

Phillips adds, "Further research could inform farmers of how to cultivate soil organic matter useful for nitrogen management. Organic forms of <u>soil nitrogen</u>, such as waste products from plants and fungi, could help convert excess inorganic nitrogen—which would otherwise be leached into water or emitted as nitrous oxide—into a form that isn't harmful to the environment."

However, the scientists say more research is needed to test exactly which forms of <u>organic nitrogen</u> are most effective. The team is now developing proposals for further funding that will allow them to investigate on-farm applications for transforming excess nitrogen from soil and water into unreactive atmospheric N2 gas without producing N2O. This may allow scientists to develop options to manage the fate of agricultural <u>nitrogen</u> while avoiding greenhouse-gas emissions.

More information: Rebecca L. Phillips et al. Chemical formation of hybrid di-nitrogen calls fungal codenitrification into question, *Scientific Reports* (2016). DOI: 10.1038/srep39077

Provided by The College of William & Mary

Citation: Finding may offer farmers a way to reduce harmful emissions from fertilized soil (2017, January 16) retrieved 4 May 2024 from <u>https://phys.org/news/2017-01-farmers-emissions-fertilized-soil.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.