


Both push and pull drive our galaxy's race through space

January 30 2017



Although we can't feel it, we're in constant motion: the earth spins on its axis at about 1,600 km/h; it orbits around the sun at about 100,000 km/h; the sun orbits our Milky Way galaxy at about 850,000 km/h; and the Milky Way galaxy and its companion galaxy Andromeda are moving with respect to the expanding universe at roughly 2 million km/h (630 km per second). But what is propelling the Milky Way's race through

space?

Until now, scientists assumed that a dense region of the universe is pulling us toward it, in the same way that gravity made Newton's apple fall to earth. The initial "prime suspect" was called the Great Attractor, a region of a half dozen rich clusters of galaxies 150 million lightyears from the Milky Way. Soon after, attention was drawn to an area of more than two dozen rich clusters, called the Shapley Concentration, which sits 600 million lightyears beyond the Great Attractor.

Now researchers led by Prof. Yehuda Hoffman at the Hebrew University of Jerusalem report that our galaxy is not only being pulled, but also pushed. In a new study in the forthcoming issue of *Nature Astronomy*, they describe a previously unknown, very large region in our extragalactic neighborhood. Largely devoid of galaxies, this void exerts a repelling force on our Local Group of galaxies.

"By 3-d mapping the flow of galaxies through space, we found that our Milky Way galaxy is speeding away from a large, previously unidentified region of [low density](#). Because it repels rather than attracts, we call this region the Dipole Repeller," said Prof. Yehuda Hoffman. "In addition to being pulled towards the known Shapley Concentration, we are also being pushed away from the newly discovered Dipole Repeller. Thus it has become apparent that push and pull are of comparable importance at our location."

The presence of such a low density region has been suggested previously, but confirming the absence of galaxies by observation has proved challenging. But in this new study, Hoffman, at the Hebrew university's Racah Institutes of Physics, working with colleagues in the USA and France, tried a different approach.

Using powerful telescopes, among them the Hubble Space Telescope,

they constructed a 3-dimensional map of the galaxy flow field. Flows are direct responses to the distribution of matter, away from regions that are relatively empty and toward regions of mass concentration; the large scale structure of the universe is encoded in the flow field of galaxies. They studied the peculiar velocities - those in excess of the Universe's rate of expansion - of galaxies around the Milky Way, combining different datasets of peculiar velocities with a rigorous statistical analysis of their properties. They thereby inferred the underlying mass distribution that consists of dark matter and luminous galaxies—over-dense regions that attract and under-dense ones that repel.

By identifying the Dipole Repeller, the researchers were able to reconcile both the direction of the Milky Way's motion and its magnitude. They expect that future ultra-sensitive surveys at optical, near-infrared and radio wavelengths will directly identify the few [galaxies](#) expected to lie in this void, and directly confirm the void associated with the Dipole Repeller.

More information: The dipole repeller, *Nature Astronomy*, [nature.com/articles/doi:10.1038/s41550-016-0036](https://doi.org/10.1038/s41550-016-0036)

Provided by Hebrew University of Jerusalem

Citation: Both push and pull drive our galaxy's race through space (2017, January 30) retrieved 27 April 2024 from <https://phys.org/news/2017-01-galaxy-space.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.